

10-2013

Second messenger/signal transduction pathways in major mood disorders: Moving from membrane to mechanism of action, part II: bipolar disorder

Mark Niciu

National Institutes of Health, Bethesda, MD

Dawn F. Ionescu

National Institutes of Health, Bethesda, MD

Daniel C. Matthews

National Institutes of Health, Bethesda, MD

Erica M. Richards

National Institutes of Health, Bethesda, MD

Carlos A. Zarate

George Washington University

Follow this and additional works at: http://hsrc.himmelfarb.gwu.edu/smhs_psych_facpubs

 Part of the [Mental and Social Health Commons](#), [Psychiatry Commons](#), and the [Psychiatry and Psychology Commons](#)

Recommended Citation

Niciu, M.J., Ionescu, D.F., Mathews, D.C., Richards, E.M., Zarate, C.A. (2013). Second messenger/signal transduction pathways in major mood disorders: Moving from membrane to mechanism of action, part II: bipolar disorder. *CNS Spectrums*, 18(5), 242-251.

This Journal Article is brought to you for free and open access by the Psychiatry and Behavioral Sciences at Health Sciences Research Commons. It has been accepted for inclusion in Psychiatry and Behavioral Sciences Faculty Publications by an authorized administrator of Health Sciences Research Commons. For more information, please contact hsrc@gwu.edu.

CNS Spectrums

<http://journals.cambridge.org/CNS>


Additional services for **CNS Spectrums**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Second messenger/signal transduction pathways in major mood disorders: moving from membrane to mechanism of action, part I: major depressive disorder

Mark J. Niciu, Dawn F. Ionescu, Daniel C. Mathews, Erica M. Richards and Carlos A. Zarate, Jr.

CNS Spectrums / Volume 18 / Issue 05 / October 2013, pp 231 - 241

DOI: 10.1017/S1092852913000059, Published online: 05 March 2013

Link to this article: http://journals.cambridge.org/abstract_S1092852913000059

How to cite this article:

Mark J. Niciu, Dawn F. Ionescu, Daniel C. Mathews, Erica M. Richards and Carlos A. Zarate, Jr. (2013). Second messenger/signal transduction pathways in major mood disorders: moving from membrane to mechanism of action, part I: major depressive disorder. *CNS Spectrums*, 18, pp 231-241 doi:10.1017/S1092852913000059

Request Permissions : [Click here](#)

Second messenger/signal transduction pathways in major mood disorders: moving from membrane to mechanism of action, part I: major depressive disorder

Mark J. Niciu,^{1*} Dawn F. Ionescu,¹ Daniel C. Mathews,¹ Erica M. Richards,¹ and Carlos A. Zarate Jr.^{1,2}

¹ National Institutes of Health (NIH)/National Institute of Mental Health (NIMH), Experimental Therapeutics and Pathophysiology Branch (ETPB), Intramural Research Program, Bethesda, Maryland, USA

² Psychiatry and Behavioral Sciences, The George Washington University

The etiopathogenesis and treatment of major mood disorders have historically focused on modulation of monoaminergic (serotonin, norepinephrine, dopamine) and amino acid [γ -aminobutyric acid (GABA), glutamate] receptors at the plasma membrane. Although the activation and inhibition of these receptors acutely alter local neurotransmitter levels, their neuropsychiatric effects are not immediately observed. This time lag implicates intracellular neuroplasticity as primary in the mechanism of action of antidepressants and mood stabilizers. The modulation of intracellular second messenger/signal transduction cascades affects neurotrophic pathways that are both necessary and sufficient for monoaminergic and amino acid-based treatments. In this review, we will discuss the evidence in support of intracellular mediators in the pathophysiology and treatment of preclinical models of despair and major depressive disorder (MDD). More specifically, we will focus on the following pathways: cAMP/PKA/CREB, neurotrophin-mediated (MAPK and others), p11, Wnt/Fz/Dvl/GSK3 β , and NF κ B/ Δ FosB. We will also discuss recent discoveries with rapidly acting antidepressants, which activate the mammalian target of rapamycin (mTOR) and release of inhibition on local translation via elongation factor stimulation. Throughout this discourse, we will highlight potential intracellular targets for therapeutic intervention. Finally, future clinical implications are discussed.

Received 7 December 2012; Accepted 11 January 2012; First published online 5 March 2013

Key words: depression, major depressive disorder, signal transduction, second messenger, intracellular cascades, antidepressants.

Clinical Implications

- Preclinical models of despair and clinical samples of major depressive disorder (MDD) reveal abnormalities in intracellular second messenger/signal transduction cascades. Some of these cascades include the following: cAMP/PKA/CREB, neurotrophin-mediated (MAPK and others), p11, Wnt/Fz/Dvl/GSK3(beta), and NF-(kappa)B/(delta)fobs.
- Deficiencies in intracellular second messenger/signal transduction pathways reverse in response to

successful treatment with traditional (monoaminergic) antidepressants.

- The rapidly-acting antidepressant ketamine induce changes in alternative intracellular cascades, e.g. mTOR activation and release of translational inhibition, in dendritic spines. These cascades are believed to be stimulated through acute antagonism of NMDA receptor and a synaptic glutamate surge.
- Intracellular second messenger/signal transduction abnormalities and reversal with successful treatment may serve as nosologica endophenotypes and biomarkers of response, respectively, to improve diagnosis and facilitate antidepressant drug development among the heterogeneity inherent in MDD.

The authors gratefully acknowledge the support of the Intramural Research Program of the National Institute of Mental Health, National Institutes of Health (IRP-NIMH/NIH, Bethesda, MD, USA), and thank the 7SE Inpatient Mood and Anxiety Disorders Research Unit of the NIMH/NIH for their support. The NIMH/NIH had no further role in the writing of this review, or in the decision to submit the paper for publication.

*Address for correspondence: Dr. Mark J. Niciu, National Institutes of Health(NIH)/National Institute of Mental Health(NIMH), Experimental Therapeutics and Pathophysiology Branch(ETPB), Intramural Research Program, 10 Center Dr., Building 10/CRC, Room 7-5545, Bethesda, MD 20814-9692, USA.

(Email: mark.niciu@nih.gov)

Introduction

The etiopathogenesis and treatment of the major mood disorders, major depressive disorder (MDD) and bipolar disorder (BD), have historically focused on the manipulation of monoaminergic (serotonin, norepinephrine, dopamine) and amino acid (γ -aminobutyric acid,

glutamate) neurotransmitters via the activation or inhibition of plasma membrane receptors. Albeit there are acute changes in local neurotransmitter levels in brain regions implicated in the pathophysiology of depression (cortex, hippocampus), antidepressant effects often require weeks to months. As a result, the “neurotransmitter imbalance” hypothesis of depression is at best incomplete. As will be displayed below, these medications ultimately elicit their effects through the activation/inhibition of intracellular signal transduction cascades. Additionally, more direct targeting of salient second messenger/signal transduction intermediates may provide more rapid and robust acting antidepressant effects than our cadre of currently available antidepressants. Also, by directly targeting these mediators, some off-target adverse events, eg, increasing serotonin levels in the gastrointestinal tract leading to dyspepsia or diarrhea, may be avoided.

In this first of two articles, we will review intracellular-mediated neuroplasticity in the pathophysiology of preclinical models of depressive-like behavior and MDD. Throughout, we will discuss progress-to-date on pathway manipulation in treatment, and at the end we will offer exciting areas for future pathophysiological studies and experimental therapeutics targeted at these intracellular neuromodulatory cascades.

Intracellular Second Messenger/Signal Transduction Cascades

cAMP/PKA/CREB

As mentioned, the delayed efficacy of monoaminergic antidepressants suggests a mechanism of action that is not simply explained by a restoration of a “chemical imbalance” via reuptake inhibition. As early as the 1980s, several preclinical investigators examined the vital role of intracellular second messenger/signal transduction cascades in the pathophysiology and treatment of depression. This examination was led by the discovery that antidepressants elicit their intracellular effects through canonical second messenger systems. Elevated synaptic levels of serotonin and norepinephrine activate cognate postsynaptic seven-transmembrane G-protein coupled receptors. Norepinephrine-induced β_1 AR and β_2 AR and serotonin-induced 5-HT₄, 5-HT₆, and 5-HT₇ receptor activation are predominantly implicated (Figure 1).^{1,2} The intracellular domain of G-protein coupled receptors interacts with G_{s/i}, which, through their α subunit, stimulates/inhibits adenylyl cyclase (AC). AC converts ATP-to-cAMP, which activates protein kinase A (PKA).

AC activity is increased with both chronic antidepressant treatment³ and electroconvulsive seizures (ECS).⁴

PKA phosphorylates downstream effector proteins involved in cytoskeletal reorganization and transcription. Standard antidepressants also increase PKA activity in fractionated rat necortex.^{4,5} The cAMP-response element binding (CREB) protein is the major transcription factor responsible for neurotrophic/protective mRNA transcription in this cascade. Like PKA, chronic antidepressants increase CREB mRNA and protein levels in the rat hippocampus.⁶ This increases expression of brain-derived neurotrophic factor (BDNF), especially in the hippocampal dentate gyrus.⁷ Transgenic CREB overexpression in the hippocampus has antidepressant-like effects in rodent models of despair, and phospho-CREB (the activated isoform) stimulates CRE-responsive gene expression with chronic antidepressant treatment.⁸

Due to aberrancies corrected by standard antidepressants, phosphodiesterase (PDE) dysfunction has been investigated in MDD. There are numerous PDE isoforms that have variable specificity for cAMP and cGMP; PDE4 is a brain-specific, cAMP-selective isoform that has been the most extensively studied in depression.⁹ As displayed by [¹¹C]-rolipram positron emission tomography (PET), PDE4 levels are globally decreased (about 20% reduction in MDD).¹⁰ As a result, PDE inhibitors have been proposed as rational therapeutic targets. An inhibitor of PDE4, (RS)-4-[3-(cyclopentyl-4-methoxy-phenyl)-2-pyrrolidin-2-one (rolipram), has antidepressant effects in both MDD¹¹ and rodent models of despair.^{12,13} Our group is currently studying changes in PDE4 levels after a treatment course with the selective serotonin reuptake inhibitor (SSRI) citalopram as a potential biomarker of treatment response (Clinical-Trials.gov identifier: NCT00369798). Even though there have been no additional trials with rolipram for two decades due to severe nausea, several pharmaceutical companies have subtype-specific, better-tolerated PDE inhibitors in their armamentarium for potential testing as antidepressants.

Neurotrophins

Centrally acting neurotrophins bind cognate receptors and intracellularly activate their tyrosine kinase domain, which induces autophosphorylation and recruits adapter proteins (Figure 1). In one of the most well-studied intracellular cascades in neuroscience, BDNF binds to TrkB, which activates the following three cascades: (1) extracellular regulated kinase (ERK)/mitogen-activated protein kinase (MAPK), (2) phospholipase C γ (PLC γ)/inositol triphosphate (IP₃), and (3) phosphatidylinositol-3 kinase (PI3K) (as shown in Figure 1, except the PLC γ /IP₃ cascade). In the ERK/MAPK cascade, TrkB autoactivation recruits several adapter proteins: Shc, Grb2, and Sos. Sos is a guanine nucleotide exchange factor that converts GDP into the more




Figure 1. Canonical signal transduction cascades in preclinical models of despair and major depressive disorder. On the left side of the figure, monoamine neurotransmitter binding to cognate receptors recruits G protein adapters to their intracellular C-terminal tail. This activates (G_s)/inhibits (G_i) the cAMP/PKA/CREB cascade. Adenylyl cyclase converts ATP-to-cAMP, which stimulates protein kinase A to phosphorylate CREB. Phospho-CREB translocates to the nucleus to stimulate transcription of target genes involved in neuroprotection, neurotransmission, and cytoskeletal dynamics. On the right, neurotrophins bind to their cognate receptor tyrosine kinases and induce the autophosphorylation of their intracellular domain(s). This recruits numerous adapter proteins to the plasma membrane and stimulates protein–protein interactions that culminate in the activation of Raf, a protein kinase. Raf activates the small molecule Ras to induce mitogen-activated protein kinase (MAPK). Like the cAMP/PKA/CREB cascade, the MAPK/ERK pathway culminates in nuclear translocation of transcription factors (including CREB) to the nucleus. There is also cross-talk between these two cascades (as depicted in the middle of the figure) via phosphoinositides leading to Akt activation. Akt phosphorylates GSK-3 β , which dissociates it from Axin and APC (“degradation complex”). This stabilizes β -catenin and facilitates its nuclear translocation. Please refer to the accompanying text for a discussion of intracellular second messenger/signal transduction aberrations in depression, normalizing responses with antidepressants, and experimental targets for future drug development. β AR, beta-adrenergic receptor; 5-HT, 5-hydroxytryptamine (serotonin); DR, dopamine receptor; Trk, tyrosine kinase; NT, neurotrophin; Flk [VEGF (vascular endothelial growth factor) receptor], fetal liver kinase; AC, adenylyl cyclase; ATP, adenosine triphosphate; cAMP, cyclic adenosine monophosphate; PDE, phosphodiesterase; CREB, cAMP-response element binding protein; LEF/TCF, lymphoid enhancer factor/T-cell factor; PP2A, protein phosphatase 2A; β -Arr, beta-arrestin; GSK-3 β , glycogen synthase kinase-3 beta; APC, adenosis polyposis coli; β -Cat, beta-catenin; SHC, Src homology-2 domain containing (protein); Grb2, growth factor receptor-bound (protein) 2; Sos, son of sevenless; Ras, rat sarcoma; GTP, guanine triphosphate; GDP, guanine diphosphate; MAPKK, mitogen-activated protein kinase kinase; ERK, extracellular-regulated kinase.

energetically rich GTP. This sequentially activates Ras and Raf, the first protein kinase in this cascade. Like PKA, the activation of ERK/MAPK stimulates the transcription of target genes responsible for cytoskeletal rearrangement, neurotransmitter secretion, reuptake, etc.

The most extensively studied neurotrophin in depression is BDNF (reviewed by Tanis and Duman¹⁴). In brief, multiple stress-induction paradigms decrease hippocampal BDNF expression and cause depression-like behaviors.¹⁵ Antidepressant medications and electroconvulsive seizures (ECS) restore BDNF mRNA levels in the frontal cortex and hippocampus.^{16,17} Exogenous administration of BDNF into the midbrain¹⁸

and hippocampus¹⁹ also has antidepressant-like effects. Conditional BDNF knockout in the mouse forebrain impairs the antidepressant effects of desipramine on the forced swim test (FST).²⁰ As a result, CNS BDNF expression is both necessary (based on inducible knockout experiments) and sufficient (from exogenous administration experiments) for antidepressant efficacy.

Vascular endothelial growth factor (VEGF) is another neurotrophin that has been investigated in depression. Stress decreases hippocampal VEGF levels.²¹ ECS restores VEGF expression and intracellular flux through its cognate receptor, Flk-1 (VEGFR2), via the proliferation of neural stem cells in the dentate gyrus²² and

recruitment of constituents of the mTORC1 signaling pathway.²³ VEGF/Flk-1 signaling is also essential for the antidepressant effects of the SSRI fluoxetine.²⁴ Next, exercise-induced alleviation of depression-like symptoms in rodents also activates VEGF/Flk-1.²³ In a rodent transgenic system (upregulation of cAMP through an *Aplysia* G_s-coupled receptor), VEGF was necessary for an antidepressant-like effect in several stress-induction paradigms.²⁵ In clinical studies, low plasma VEGF levels have been observed in suicide completers,²⁶ and the antidepressant effects of total sleep deprivation coincide with increased plasma levels of VEGF.²⁷ In a combined cohort of subjects in a current major depressive episode (both unipolar and bipolar depression), higher pretreatment VEGF levels trended in antidepressant responders versus nonresponders ($p = 0.055$).²⁸ Peripheral VEGF levels also remained elevated up to 1 month after a successful course of ECT.²⁹ On a genetic level, the VEGF C/A polymorphism is associated with treatment-resistant depression (TRD), as the CC genotype is more common in ECT-treated patients than in controls (31.1% and 18.7%, respectively).³⁰ However, another pharmacogenetic study revealed no association of seven different VEGF polymorphisms and antidepressant response.³¹

Insulin-like growth factor-1 (IGF-1) has also been studied in depression. IGF-1 is produced by neuroendocrine cells in response to circulating hormones, especially growth hormone (GH) and insulin, and has potent mitogenic effects.³² Although initial clinical reports demonstrated increased IGF-1 levels in depressed patients,^{33,34} these investigations only examined peripheral levels (which may not accurately reflect centrally acting IGF-1) and did not discern between free and bound IGF-1.³⁵ As a result, several preclinical research groups have clarified the role of centrally acting IGF-1 and its inhibition. IGF-1 knockdown in CA1 hippocampal pyramidal neurons has depressogenic effects.³⁶ Intracerebroventricular administration of IGF-1 and a non-selective IGF binding protein (which sequesters IGF-1 into a biologically inert complex) inhibitor have antidepressant and anxiolytic-like effects in stress induction paradigms.³⁷ Central IGF-1 also decreases expression of proinflammatory cytokines, which may mitigate neuroinflammatory cascades that are critical in depression onset and/or maintenance.^{38,39} Peripheral administration of an IGF-1 antibody blocks the antidepressant effects of exercise in a murine model of chronic unpredictable stress.⁴⁰ Back in the clinic, antidepressant treatment increased low CSF levels of IGF-1⁴¹; as a result, exogenous immediately acting (intranasal) IGF-1 is being investigated for the treatment of MDD.⁴²

Due to the discovery of decreased glial cell numbers in rodent models of despair and MDD,⁴³ several groups have investigated a putative role for glial-

derived neurotrophic factor (GDNF) in depression. MDD patients display an age-dependent decrease in peripheral GDNF levels,^{44,45} which increases in response to treatment^{46,47} and normalizes during remission.⁴⁸ In contrast, in a postmortem sample of recurrent depression, increased GDNF levels in parietal cortex were evident. Finally, in a study of rat glioma cells, antidepressant-induced GDNF expression/secretion was mediated by β -arrestin-1/CREB transcription complex formation,⁴⁹ and GDNF epigenetic regulation (promoter methylation and histone modification) had adaptive effects in stressed mice.⁵⁰

Other centrally expressed neurotrophins, eg, nerve growth factor (NGF) and neurotrophin-3 (NT-3) and their cognate receptors TrkA, TrkC and p75NTR, are also under investigation in preclinical/clinical studies.

p11

p11 was initially found in a yeast two-hybrid screen as a 5-HT_{1B} and 5-HT₄ interactor.⁵¹ p11 mRNA and 5-HT_{1B} receptor transcripts co-express in several brain areas salient for depression.⁵² p11 mRNA was compared in helpless H/Rouen mice (a genetic model of depression) versus non-helpless NH/Rouen mice, and, at baseline, p11 mRNA levels were decreased in the forebrain in the helpless H/Rouen mice.⁵¹ The antidepressants imipramine and tranylcypromine as well as ECS increase neocortical p11 mRNA levels.⁵¹ p11 knockout mice display biochemical, electrophysiological, and behavioral responses consistent with depression.⁵¹ Furthermore, the antidepressant effects of imipramine in p11^{-/-} mice were reduced in these mice, and, when exposed to tail suspension and FST, they were more resistant to the antidepressant effects of exogenous BDNF.⁵³ The transgenic overexpression of p11, on the other hand, increases resiliency to exogenous stress. BDNF increases p11 expression in a trkB and MAPK-dependent manner.⁵³ Therefore, p11 is both necessary in the pathogenesis and sufficient for reversal of depressive behaviors.

The expression of p11 mRNA has also been examined in depressed suicide completers, which revealed downregulation in the anterior cingulate cortex relative to non-depressed controls.⁵² Prefrontal p11 mRNA is also decreased in suicide completers relative to postmortem controls.⁵⁴ Peripheral p11 mRNA levels are also decreased in suicidal attempters with comorbid MDD and posttraumatic stress disorder (PTSD) relative to non-attempters⁵⁴ (but increased relative to PTSD alone and healthy volunteers in another sample⁵⁵).

In non-human primates, chronic treatment with fluoxetine significantly increased p11 in peripheral mononuclear cells temporally consistent with antidepressant onset (unpublished personal communication




Figure 2. Canonical Wnt/Fz/Dvl/GSK-3 β signal transduction cascade. Wnts are secreted glycoproteins that are agonists for Fz receptors. Fz receptor stimulation recruits LRP to the plasma membrane, which may be inhibited by cytosolic isoforms of Dkk. The Wnt-Fz ligand-receptor complex recruits the scaffolding protein Dvl, which, in turn, stimulates Ca²⁺-dependent and Ca²⁺-independent pathways. The Ca²⁺ dependent pathway activates, among other mediators, PKC and CAMKII. In addition to their more notorious effects on phosphoinositides, PKC crosstalk with the MAPK/ERK cascade promotes neuromodulatory gene transcription (as depicted in Figure 1). CAMKII binds Ca²⁺ and stimulates neuromodulatory gene transcription through CREB. In the Ca²⁺ independent arm, Dvl stimulates phosphorylation of GSK-3 β (as described in Figure 1) and facilitates dissociation of the β -catenin degradation complex and β -catenin mediated gene transcription. Please refer to the accompanying text for a discussion of aberrations in depression, normalizing responses with successful antidepressant treatment, and potential experimental targets for future investigation. Fz, frizzled; LRP, low-density lipoprotein receptor-related protein; Dkk, Dickkopf; Dvl, disheveled; PKC, protein kinase C; MAPK, mitogen activated protein kinase; ERK, extracellular-regulated kinase; CAMKII, calcium-calmodulin dependent protein kinase II; CREB, cAMP-response element binding protein; LEF/TCF, lymphoid enhancer factor/T-cell factor; GSK-3 β , glycogen synthase kinase-3 beta; APC, adenosis polyposis coli; β -Cat, beta-catenin.

from R. Innis, MD, PhD, NIMH). To translate these findings into humans, our group is presently investigating if peripheral p11 levels increase in response to successful SSRI treatment as a potentially biologically salient biomarker of treatment response (Clinical-Trials.gov identifier: NCT00697268).

Wnt/Fz/Dvl/GSK-3 β

The Wnt/frizzled/disheveled/glycogen synthase kinase-3 beta cascade has been studied in the pathophysiology and therapeutics of depression (Figure 2). (Of note, GSK-3 β can also be activated by PI3K and Akt; for a review, see Voleti and Duman.)⁵⁶ In the canonical signal transduction cascade, Wnt binding to Fz recruits a low-density lipoprotein receptor-related protein (LRP)5/6 to the plasma membrane, which interacts with the scaffolding protein disheveled. Disheveled mediates GSK-3 β phosphorylation, which inactivates it. This releases β -catenin from the axin-adenosis polyposis

coli-GSK-3 β "destruction complex" for nuclear translocation.⁵⁷ Nuclear β -catenin interacts with the transcription factor T-cell factor/lymphoid enhancer factor (TCF/LEF) to express Wnt-responsive genes.

The Wnt/Fz/Dvl/GSK-3 β cascade has been implicated in neuromodulation, especially synapse formation, neurotransmission, and cytoskeletal reorganization.⁵⁸ The expression of an endogenous Wnt inhibitor, Dickkopf-1 (Dkk-1), is increased with mild restraint stress and exogenous corticosterone administration.⁵⁹ Mice lacking the Dkk1 transcriptional enhancer (Doubldridge mice) are more resilient to chronic unpredictable stress.⁵⁹ Another isoform, Dickkopf-2 (Dkk2), is down-regulated by chronic ECS.⁶⁰ In this same study, a frizzled receptor isoform, Fz6, was increased by chronic ECS⁶⁰ and demonstrated that viral vector-mediated inhibition of Fz6 was anxi- and depressogenic in numerous behavioral paradigms.

GSK-3 β is a serine-threonine kinase that has been extensively investigated in psychotic and mood

disorders, especially bipolar disorder after the discovery that lithium is a potent GSK-3 β inhibitor.⁶¹ In preclinical models of despair, the heterozygous deletion of GSK-3 β has antidepressant effects, and GSK-3 inhibitors (L803-*mts* and AR-A014418) mimicked these genetic effects.^{60,62,63} The phosphorylation of GSK-3 β is increased by chronic administration of the antidepressants fluoxetine or venlafaxine.⁶⁴ A depressive phenotype was also observed with overexpression of GSK3 β in the NAcc, while a dominant-negative GSK3 β isoform promoted resiliency.⁶⁵

Several other pathway intermediates are affected by antidepressant therapy. Wnt2 was increased by several antidepressants (including ECS) in a microarray study of a rodent model of despair.⁶⁴ Additionally, Wnt2 transgenic overexpression in the hippocampus was sufficient to generate antidepressant-like effects.⁶⁴ Wnt7b expression, on the other hand, was increased by atomoxetine and ECS.⁶⁴ Fz9 levels were upregulated by the noradrenergic antidepressants atomoxetine and venlafaxine but not by SSRIs.⁶⁴ Blockade of a disheveled isoform, Dvl2 [via both overexpression of a dominant-negative isoform and intra-nucleus accumens (NAcc) inhibitor infusion], decreased resiliency to social defeat and other modalities for inducing despair.⁶⁵

To our knowledge, there have been no CNS-penetrant small molecule modulators of the Wnt/Fz/Dvl/GSK-3 β pathway that have been tested in psychiatric disorders. We seek translation of these interesting rodent findings into the pathophysiology and experimental therapeutics of MDD. Due to this signal transduction/second messenger system's involvement in numerous cellular pathways, especially mitogenesis, translational studies will need to pay close attention to toxicity and side effect profiles.

NF- κ B/ Δ FosB

Nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) is a transcription factor that has been most extensively studied in immunology but also plays a functional role in synaptic processes underlying learning and memory.⁶⁶ After demonstrating its function in instrumental and other forms of motivated learning, several groups have investigated NF- κ B induction of Δ FosB, a truncated version of the immediate-early transcription factor FosB, in rodent models of despair. Lipopolysaccharide-exposed rodents have a delayed increase in immunostaining for Δ FosB that parallels the onset of increased immobility on the FST and reduced sucrose consumption (a preclinical marker of anhedonia).⁶⁷ Serum response factor (SRF), another transcription factor, decreases Δ FosB in the NAcc in response to chronic social defeat.⁶⁸ SRF levels are also

decreased in the NAcc in chronic social defeat stress in mice and unipolar depression.⁶⁸ In addition, genetic deletion of NAcc SRF decreased resiliency to stress.⁶⁸

Several studies have demonstrated that Δ FosB expression correlates with antidepressant-like effects.⁶⁹ In an elegant study using different mouse strains genetically engineered to produce FosB +/- Δ FosB, the Δ FosB haplotype (+/ Δ) strain had increased depressive-like behaviors relative to wild-type and the double knock-in (Δ/Δ), which displayed less anxiety in the open field test.⁷⁰ Δ FosB expression increased in multiple rat brain regions, eg, dorsal raphe nucleus, frontal cortex, hippocampus, and basolateral amygdala, with standard antidepressants (sertraline and desipramine) and vagal nerve stimulation (VNS).⁷¹ Interestingly, VNS appeared to have a larger effect and affected some brain regions not observed with traditional antidepressants (nucleus tractus solitarius and locus ceruleus).⁷¹

mTOR

As the activation of intracellular neuromodulatory cascades is critical for the mechanism of action of standard antidepressants, several recent studies have elucidated the mechanisms underlying the rapidly acting antidepressant effects of glutamate-based medications such as ketamine. Li *et al.*⁷² discerned that the activation of mammalian target of rapamycin (mTOR) was necessary for ketamine's antidepressant effects (Figure 3). A case report in a single treatment-resistant depressed patient revealed that intravenous ketamine increased peripheral mTOR expression on a time course that coincided with its rapid antidepressant effects.⁷³ Like ketamine, the proprietary mGluR2/3 antagonist, LY341495, rapidly (within 1 h) activated mTOR and downstream pathway constituents (p70S6K, 4E-BP1) and subsequently (24 h later) increased levels of postsynaptic density proteins (PSD-95, GluR1, synapsin I).⁷⁴ These antidepressant effects of LY341495 were reversed by the mTOR inhibitor rapamycin.⁷⁴

eEF2K/CAMKIII

Finally, the release of inhibition on local translation in dendritic spines has emerged as an exciting intracellular target of ketamine. Autry *et al.*⁷⁵ reported that subanesthetic doses of ketamine released inhibition of translation by deactivating eukaryotic elongation factor 2 kinase (eEF2K)/calcium-calmodulin protein kinase type III (CAMKIII) (Figure 3). The ensuing dephosphorylation of eEF2 removes tonic inhibition on BDNF translation in the hippocampus, thereby increasing BDNF levels and concomitant TrkB receptor activation. The deactivation of eEF2K and stimulation of eEF2




Figure 3. Signal transduction cascades activated by the rapidly-acting antidepressant ketamine. Based on preliminary preclinical and unpublished clinical data, postsynaptic NMDA receptor antagonism incites a rapid presynaptic glutamate surge. Glutamate then stimulates AMPA receptors. AMPA potentiation increases Na⁺ to induce the phosphorylation of mTOR, a central signaling hub that has multiple downstream effectors. Activated mTOR then phosphorylates p70S6K. Phospho-p70S6K can directly stimulate translation of downstream postsynaptic targets. Stimulated mTOR also inhibits 4E-BP, thereby relieving translational inhibition. NMDA receptor activation also inhibits eEF2K, which increases levels of dephosphorylated eEF2. Dephosphorylated eEF2 relieves inhibition of BDNF translation in dendritic spines. This multitiered translational activation increases the expression of neuromodulatory proteins involved in, among other effects, postsynaptic scaffolding, neurotransmitter dynamics, and dendritic spine morphogenesis (from synaptically unstable filopodia to synaptically dynamic mushroom-shaped spines, which form the morphological substrate for diverse neuropsychiatric responses: learning, memory, and, most importantly here, antidepressant-like behavioral effects). Through its release of inhibition on local translation, ketamine also increases excitatory postsynaptic potentials in prefrontal cortical neurons. NMDA, N-methyl-D-aspartate; AMPA, 2-amino-3-(3-hydroxy-5-methylisoxazol-4-yl)propanoic acid; mTOR, mammalian target of rapamycin; 4E-BP, eukaryotic initiation factor 4E-binding protein; eEF2, eukaryotic elongation factor 2; eEF2K, eEF2 kinase; FKBP12, FK506-binding protein 12; LST8, mammalian *lethal with sec 13*; PI3K, phosphoinositide-3 kinase; mTORC1, mTOR complex 1; PRAS40, proline-rich Akt/PKB substrate of 40 kD; p70S6K, p70 S6 kinase. T = threonine, S = serine (with trailing number denoting the phosphorylated residue).

(thereby increasing central BDNF levels) has emerged as a novel rational therapeutic target in MDD.⁷⁶

Conclusions

In this article, we have reviewed the evidence for intracellular second messenger/signal transduction cascades in preclinical models of despair and MDD. Monoamine reuptake inhibition by traditional antidepressants occurs immediately, but the behavioral effects take much longer, which implicates intracellular processes in their mechanism of action. Traditional monoaminergic antidepressants indirectly stimulate multiple intracellular cascades, and this may ultimately dilute their antidepressant efficacy on the neural circuitry that is involved in depression via off-target stimulation/inhibition. In addition, these

medications have proven inadequate in real world effectiveness trials such as STAR*D⁷⁷ and CO-MED.⁷⁸ This should come as no surprise, as only 5–10% of CNS neurons use monoamines as their primary neurotransmitter, while >50% use glutamate. Glutamate-based antidepressants are more rapidly acting and have larger effect sizes in clinical trials.^{79,80} Two of the most promising candidates for future drug development based on preclinical studies with ketamine are mTOR and eEF2K/eEF2. As reviewed, both of these molecules are critical in central BDNF signaling (mTOR possibly downstream of BDNF and eEF2K/eEF2 in local BDNF translation). As promising as this seems, we must remain vigilant for toxicity/adverse events, as (1) mTOR is a protooncogene and overactive in autoimmune disorders (where its overactivity may be suppressed clinically with rapamycin), and (2) chronic stimulation of eEF2 may

lead to excessive translation of off-target proteins. (Of note, we suggest that monoaminergic antidepressants have not been associated with an increased risk of cancer because their biological effects are more indirect and likely less potent on these intracellular cascades.)

The mood disorders field is sorely in need of biologically salient measures to improve our existing nosology and monitor treatment response, which, at present, is based only on patient report and clinical impression. Biologically informed therapeutics are routinely used in other fields of medicine such as cardiology and oncology, but, unfortunately, they remain elusive in psychiatry. Reliable measures of intracellular processes involved in depression may assist in developing a more accurate nosology among the heterogeneity inherent in the clinical diagnosis of MDD, eg, those patients who have underactive mTOR or eEF2 activity may have a "glutamate-based depression"⁸¹ and may benefit from glutamate-based therapies. Next, the development of peripheral measures of intracellular events may allow us to assess a more reliable and quantitative baseline and treatment response better than our current approach, eg, assessing baseline mTOR and eEF2K/eEF2 activity and monitoring change in activity with treatment.

In conclusion, we have garnered an adequate understanding of intracellular second messenger/signal transduction cascades in preclinical models of depression and MDD, and, in due time, these findings will likely be translated into the clinic in novel therapies and nosological biomarkers.

Disclosures

The authors gratefully acknowledge the support of the IRP-NIMH/NIH, and the NARSAD Independent Investigator Award and Brain and Behavior Foundation Bipolar Research Award (Dr. Zarate). Salary support was also provided by the IRP-NIMH/NIH (MJN, DFI, DCM, and EMR). Drs. Niciu, Ionescu, Mathews, and Richards have no potential financial conflicts of interest to disclose. Dr. Zarate is listed as a co-inventor on a patent application for the use of ketamine and its metabolites in major depression. Dr. Zarate has assigned his rights in the patent to the U.S. Government but will share a percentage of any royalties that may be received. Dr. Zarate is also a government employee of the NIH.

References

1. Sleight AJ, Carolo C, Petit N, Zwingelstein C, Bourson A. Identification of 5-hydroxytryptamine₇ receptor binding sites in rat hypothalamus: sensitivity to chronic antidepressant treatment. *Mol Pharmacol*. 1995; **47**(1): 99–103.
2. Svenningsson P, Tzavara ET, Witkin JM, et al. Involvement of striatal and extrastriatal DARPP-32 in

- biochemical and behavioral effects of fluoxetine (Prozac). *Proc Natl Acad Sci U S A*. 2002; **99**(5): 3182–3187.
3. Menkes DB, Rasenick MM, Wheeler MA, Bitensky MW. Guanosine triphosphate activation of brain adenylate cyclase: enhancement by long-term antidepressant treatment. *Science*. 1983; **219**(4580): 65–67.
4. Ozawa H, Rasenick MM. Chronic electroconvulsive treatment augments coupling of the GTP-binding protein Gs to the catalytic moiety of adenylyl cyclase in a manner similar to that seen with chronic antidepressant drugs. *J Neurochem*. 1991; **56**(1): 330–338.
5. Nestler EJ, Terwilliger RZ, Duman RS. Chronic antidepressant administration alters the subcellular distribution of cyclic AMP-dependent protein kinase in rat frontal cortex. *J Neurochem*. 1989; **53**(5): 1644–1647.
6. Nibuya M, Nestler EJ, Duman RS. Chronic antidepressant administration increases the expression of cAMP response element binding protein (CREB) in rat hippocampus. *J Neurosci*. 1996; **16**(7): 2365–2372.
7. Conti AC, Cryan JF, Dalvi A, Lucki I, Blendy JA. cAMP response element-binding protein is essential for the upregulation of brain-derived neurotrophic factor transcription, but not the behavioral or endocrine responses to antidepressant drugs. *J Neurosci*. 2002; **22**(8): 3262–3268.
8. Thome J, Sakai N, Shin K, et al. cAMP response element-mediated gene transcription is upregulated by chronic antidepressant treatment. *J Neurosci*. 2000; **20**(11): 4030–4036.
9. Zhang HT. Cyclic AMP-specific phosphodiesterase-4 as a target for the development of antidepressant drugs. *Curr Pharm Des*. 2009; **15**(14): 1688–1698.
10. Fujita M, Hines CS, Zoghbi SS, et al. Downregulation of brain phosphodiesterase type IV measured with (11)C-(R)-rolipram positron emission tomography in major depressive disorder. *Biol Psychiatry*. 2012; **72**(7): 548–554.
11. Fleischhacker WW, Hinterhuber H, Bauer H, et al. A multicenter double-blind study of three different doses of the new cAMP-phosphodiesterase inhibitor rolipram in patients with major depressive disorder. *Neuropsychobiology*. 1992; **26**(1–2): 59–64.
12. Fujimaki K, Morinobu S, Duman RS. Administration of a cAMP phosphodiesterase 4 inhibitor enhances antidepressant-induced of BDNF mRNA in rat hippocampus. *Neuropsychopharmacology*. 2000; **22**(1): 42–51.
13. Itoh T, Tokumura M, Abe K. Effects of rolipram, a phosphodiesterase 4 inhibitor, in combination with imipramine on depressive behavior, CRE-binding activity and BDNF level in learned helplessness rats. *Eur J Pharmacol*. 2004; **498**(1–3): 135–142.
14. Tanis KQ, Duman RS. Intracellular signaling pathways pave roads to recovery for mood disorders. *Ann Med*. 2007; **39**(7): 531–544.
15. Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry*. 2006; **59**(12): 1116–1127.
16. Nibuya M, Morinobu S, Duman RS. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J Neurosci*. 1995; **15**(11): 7539–7547.

17. Russo-Neustadt A, Beard RC, Cotman CW. Exercise, antidepressant medications, and enhanced brain derived neurotrophic factor expression. *Neuropsychopharmacology*. 1999; **21**(5): 679–682.
18. Siuciak JA, Lewis DR, Wiegand SJ, Lindsay RM. Antidepressant-like effect of brain-derived neurotrophic factor (BDNF). *Pharmacol Biochem Behav*. 1997; **56**(1): 131–137.
19. Shirayama Y, Chen AC, Nakagawa S, Russell DS, Duman RS. Brain-derived neurotrophic factor produces antidepressant effects in behavioral models of depression. *J Neurosci*. 2002; **22**(8): 3251–3261.
20. Monteggia LM, Barrot M, Powell CM, et al. Essential role of brain-derived neurotrophic factor in adult hippocampal function. *Proc Natl Acad Sci U S A*. 2004; **101**(29): 10827–10832.
21. Heine VM, Zareno J, Maslam S, Joels M, Lucassen PJ. Chronic stress in the adult dentate gyrus reduces cell proliferation near the vasculature and VEGF and Flk-1 protein expression. *Eur J Neurosci*. 2005; **21**(5): 1304–1314.
22. Segi-Nishida E, Warner-Schmidt JL, Duman RS. Electroconvulsive seizure and VEGF increase the proliferation of neural stem-like cells in rat hippocampus. *Proc Natl Acad Sci U S A*. 2008; **105**(32): 11352–11357.
23. Elfving B, Wegener G. Electroconvulsive seizures stimulate the VEGF pathway via mTORC1. *Synapse*. 2012; **66**(4): 340–345.
24. Greene J, Banasr M, Lee B, Warner-Schmidt J, Duman RS. Vascular endothelial growth factor signaling is required for the behavioral actions of antidepressant treatment: pharmacological and cellular characterization. *Neuropsychopharmacology*. 2009; **34**(11): 2459–2468.
25. Lee JS, Jang DJ, Lee N, et al. Induction of neuronal vascular endothelial growth factor expression by cAMP in the dentate gyrus of the hippocampus is required for antidepressant-like behaviors. *J Neurosci*. 2009; **29**(26): 8493–8505.
26. Isung J, Mobarrez F, Nordstrom P, Asberg M, Jokinen J. Low plasma vascular endothelial growth factor (VEGF) associated with completed suicide. *World J Biol Psychiatry*. 2012; **13**(6): 468–473.
27. Ibrahim L, Duncan W, Luckenbaugh DA, et al. Rapid antidepressant changes with sleep deprivation in major depressive disorder are associated with changes in vascular endothelial growth factor (VEGF): a pilot study. *Brain Res Bull*. 2011; **86**(1–2): 129–133.
28. Halmai Z, Dome P, Dobos J, et al. Peripheral vascular endothelial growth factor level is associated with antidepressant treatment response: results of a preliminary study. *J Affect Disord*. 2013; **144**(3): 269–273.
29. Minelli A, Zanardini R, Abate M, et al. Vascular endothelial growth factor (VEGF) serum concentration during electroconvulsive therapy (ECT) in treatment resistant depressed patients. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; **35**(5): 1322–1325.
30. Viikki M, Anttila S, Kampman O, et al. Vascular endothelial growth factor (VEGF) polymorphism is associated with treatment resistant depression. *Neurosci Lett*. 2010; **477**(3): 105–108.
31. Tsai SJ, Hong CJ, Liou YJ, et al. Haplotype analysis of single nucleotide polymorphisms in the vascular endothelial growth factor (VEGFA) gene and antidepressant treatment response in major depressive disorder. *Psychiatry Res*. 2009; **169**(2): 113–117.
32. Clemmons DR. Modifying IGF1 activity: an approach to treat endocrine disorders, atherosclerosis and cancer. *Nat Rev Drug Discov*. 2007; **6**(10): 821–833.
33. Lesch KP, Rupperecht R, Muller U, Pfuller H, Beckmann H. Insulin-like growth factor I in depressed patients and controls. *Acta Psychiatr Scand*. 1988; **78**(6): 684–688.
34. Deuschle M, Blum WF, Strasburger CJ, et al. Insulin-like growth factor-I (IGF-I) plasma concentrations are increased in depressed patients. *Psychoneuroendocrinology*. 1997; **22**(7): 493–503.
35. Weber-Hamann B, Blum WF, Kratzsch J, et al. Insulin-like growth factor-I (IGF-I) serum concentrations in depressed patients: relationship to saliva cortisol and changes during antidepressant treatment. *Pharmacopsychiatry*. 2009; **42**(1): 23–28.
36. Mitschelen M, Yan H, Farley JA, et al. Long-term deficiency of circulating and hippocampal insulin-like growth factor I induces depressive behavior in adult mice: a potential model of geriatric depression. *Neuroscience*. 2011; **185**: 50–60.
37. Malberg JE, Platt B, Rizzo SJ, et al. Increasing the levels of insulin-like growth factor-I by an IGF binding protein inhibitor produces anxiolytic and antidepressant-like effects. *Neuropsychopharmacology*. 2007; **32**(11): 2360–2368.
38. Park SE, Dantzer R, Kelley KW, McCusker RH. Central administration of insulin-like growth factor-I decreases depressive-like behavior and brain cytokine expression in mice. *Journal of Neuroinflammation*. 2011; **8**: 12.
39. Park SE, Lawson M, Dantzer R, Kelley KW, McCusker RH. Insulin-like growth factor-I peptides act centrally to decrease depression-like behavior of mice treated intraperitoneally with lipopolysaccharide. *Journal of Neuroinflammation*. 2011; **8**: 179.
40. Duman CH, Schlesinger L, Terwilliger R, et al. Peripheral insulin-like growth factor-I produces antidepressant-like behavior and contributes to the effect of exercise. *Behav Brain Res*. 2009; **198**(2): 366–371.
41. Schilling C, Blum WF, Heuser I, et al. Treatment with antidepressants increases insulin-like growth factor-I in cerebrospinal fluid. *J Clin Psychopharmacol*. 2011; **31**(3): 390–392.
42. Paslakis G, Blum WF, Deuschle M. Intranasal insulin-like growth factor I (IGF-I) as a plausible future treatment of depression. *Med Hypotheses*. 2012; **79**(2): 222–225.
43. Rajkowska G, Miguel-Hidalgo JJ. Gliogenesis and glial pathology in depression. *CNS Neurol Disord Drug Targets*. 2007; **6**(3): 219–233.

44. Diniz BS, Teixeira AL, Miranda AS, et al. Circulating glial-derived neurotrophic factor is reduced in late-life depression. *J Psychiatr Res.* 2012; **46**(1): 135–139.
45. Tseng PT, Lee Y, Lin PY. Age-associated decrease in serum glial cell line-derived neurotrophic factor levels in patients with major depressive disorder. *Progr Neuropsychopharmacol Biol Psychiatry.* 2013; **40**: 334–339.
46. Zhang X, Zhang Z, Sha W, et al. Electroconvulsive therapy increases glial cell-line derived neurotrophic factor (GDNF) serum levels in patients with drug-resistant depression. *Psychiatry Res.* 2009; **170**(2–3): 273–275.
47. Liu Q, Zhu HY, Li B, et al. Chronic clomipramine treatment restores hippocampal expression of glial cell line-derived neurotrophic factor in a rat model of depression. *J Affect Disord.* 2012; **141**(2–3): 367–372.
48. Otsuki K, Uchida S, Watanuki T, et al. Altered expression of neurotrophic factors in patients with major depression. *J Psychiatr Res.* 2008; **42**(14): 1145–1153.
49. Golan M, Schreiber G, Avissar S. Antidepressants elevate GDNF expression and release from C(6) glioma cells in a beta-arrestin1-dependent, CREB interactive pathway. *Int J Neuropsychopharmacol.* 2011; **14**(10): 1289–1300.
50. Uchida S, Hara K, Kobayashi A, et al. Epigenetic status of GDNF in the ventral striatum determines susceptibility and adaptation to daily stressful events. *Neuron.* 2011; **69**(2): 359–372.
51. Svenningsson P, Chergui K, Rachleff I, et al. Alterations in 5-HT1B receptor function by p11 in depression-like states. *Science.* 2006; **311**(5757): 77–80.
52. Anisman H, Du L, Palkovits M, et al. Serotonin receptor subtype and p11 mRNA expression in stress-relevant brain regions of suicide and control subjects. *J Psychiatry Neurosci.* 2008; **33**(2): 131–141.
53. Warner-Schmidt JL, Chen EY, Zhang X, et al. A role for p11 in the antidepressant action of brain-derived neurotrophic factor. *Biol Psychiatry.* 2010; **68**(6): 528–535.
54. Zhang L, Su TP, Choi K, et al. P11 (S100A10) as a potential biomarker of psychiatric patients at risk of suicide. *J Psychiatr Res.* 2011; **45**(4): 435–441.
55. Su TP, Zhang L, Chung MY, et al. Levels of the potential biomarker p11 in peripheral blood cells distinguish patients with PTSD from those with other major psychiatric disorders. *J Psychiatr Res.* 2009; **43**(13): 1078–1085.
56. Voleti B, Duman RS. The roles of neurotrophic factor and Wnt signaling in depression. *Clin Pharmacol Ther.* 2012; **91**(2): 333–338.
57. Stamos JL, Weis WI. The β -catenin destruction complex. *Cold Spring Harb Perspect Biol.* 2013; **5**(1). <http://www.ncbi.nlm.nih.gov/pubmed/23169527>.
58. Inestrosa NC, Arenas E. Emerging roles of Wnts in the adult nervous system. *Nat Rev Neurosci.* 2010; **11**: 77–86.
59. Matrisciano F, Busceti CL, Bucci D, et al. Induction of the Wnt antagonist Dickkopf-1 is involved in stress-induced hippocampal damage. *PLoS One.* 2011; **6**(1): e16447.
60. Voleti B, Tanis KQ, Newton SS, Duman RS. Analysis of target genes regulated by chronic electroconvulsive therapy reveals role for Fzd6 in depression. *Biol Psychiatry.* 2012; **71**(1): 51–58.
61. Machado-Vieira R, Manji HK, Zarate CA Jr. The role of lithium in the treatment of bipolar disorder: convergent evidence for neurotrophic effects as a unifying hypothesis. *Bipolar Disord.* 2009; **11**(suppl 2): 92–109.
62. Gould TD, Einat H, Bhat R, Manji HK. AR-A014418, a selective GSK-3 inhibitor, produces antidepressant-like effects in the forced swim test. *Int J Neuropsychopharmacol.* 2004; **7**(4): 387–390.
63. O'Brien WT, Harper AD, Jove F, et al. Glycogen synthase kinase-3 β haploinsufficiency mimics the behavioral and molecular effects of lithium. *J Neurosci.* 2004; **24**(30): 6791–6798.
64. Okamoto H, Voleti B, Banasr M, et al. Wnt2 expression and signaling is increased by different classes of antidepressant treatments. *Biol Psychiatry.* 2010; **68**(6): 521–527.
65. Wilkinson MB, Dias C, Magida J, et al. A novel role of the WNT-dishevelled-GSK3 β signaling cascade in the mouse nucleus accumbens in a social defeat model of depression. *J Neurosci.* 2011; **31**: 9084–9092.
66. Meffert MK, Baltimore D. Physiological functions for brain NF-kappaB. *Trends Neurosci.* 2005; **28**(1): 37–43.
67. Frenois F, Moreau M, O'Connor J, et al. Lipopolysaccharide induces delayed FosB/DeltaFosB immunostaining within the mouse extended amygdala, hippocampus and hypothalamus, that parallel the expression of depressive-like behavior. *Psychoneuroendocrinology.* 2007; **32**(5): 516–531.
68. Vialou V, Maze I, Renthal W, et al. Serum response factor promotes resilience to chronic social stress through the induction of DeltaFosB. *J Neurosci.* 2010; **30**(43): 14585–14592.
69. McClung CA, Ulery PG, Perrotti LI, et al. DeltaFosB: a molecular switch for long-term adaptation in the brain. *Brain Res Mol Brain Res.* 2004; **132**(2): 146–154.
70. Ohnishi YN, Ohnishi YH, Hokama M, et al. FosB is essential for the enhancement of stress tolerance and antagonizes locomotor sensitization by DeltaFosB. *Biol Psychiatry.* 2011; **70**(5): 487–495.
71. Furmaga H, Sadhu M, Frazer A. Comparison of DeltaFosB immunoreactivity induced by vagal nerve stimulation with that caused by pharmacologically diverse antidepressants. *J Pharmacol Exp Ther.* 2012; **341**(12): 317–325.
72. Li N, Lee B, Liu RJ, et al. mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science.* 2010; **329**(5994): 959–964.
73. Denk MC, Rewerts C, Holsboer F, Erhardt-Lehmann A, Turck CW. Monitoring ketamine treatment response in a depressed patient via peripheral mammalian target of rapamycin activation. *Am J Psychiatry.* 2011; **168**(7): 751–752.
74. Dwyer JM, Lepack AE, Duman RS. mTOR activation is required for the antidepressant effects of mGluR(2)/(3) blockade. *Int J Neuropsychopharmacol.* 2012; **15**(4): 429–434.

75. Autry AE, Adachi M, Nosyreva E, *et al.* NMDA receptor blockade at rest triggers rapid behavioural antidepressant responses. *Nature*. 2011; **475**(7354): 91–95.
76. Monteggia LM, Gideons E, Kavalali ET. The role of eukaryotic elongation factor 2 kinase in rapid antidepressant action of ketamine. *Biol Psychiatry* In press. DOI: 10.1016/j.biopsych.2012.09.006.
77. Rush AJ, Trivedi MH, Wisniewski SR, *et al.* Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry*. 2006; **163**(11): 1905–1917.
78. Rush AJ, Trivedi MH, Stewart JW, *et al.* Combining medications to enhance depression outcomes (CO-MED): acute and long-term outcomes of a single-blind randomized study. *Am J Psychiatry*. 2011; **168**(7): 689–701.
79. Zarate CA Jr, Singh JB, Carlson PJ, *et al.* A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch Gen Psychiatry*. 2006; **63**(8): 856–864.
80. Zarate CA Jr, Mathews D, Ibrahim L, *et al.* A randomized trial of a low-trapping nonselective n-methyl-d-aspartate channel blocker in major depression. *Biol Psychiatry* In press. DOI: 10.1016/j.biopsych.2012.10.019.
81. McCarthy DJ, Alexander R, Smith MA, *et al.* Glutamate-based depression GBD. *Med Hypotheses*. 2012; **78**(5): 675–681.