

11-15-2013

The prince and the pauper: Which one is real?: The problem of secondary mutation during mutagenesis in *Streptococcus pyogenes*

Kyu Hong Cho
Southern Illinois University

Jordan O. Wright
Southern Illinois University

Juan P. Svencionis
Southern Illinois University

Jeong-Ho Kim
George Washington University

Follow this and additional works at: http://hsrc.himmelfarb.gwu.edu/smhs_biochem_facpubs

 Part of the [Microbiology Commons](#)

Recommended Citation

Cho, K., Wright, J., Svencionis, J., & Kim, J. (2013). The prince and the pauper: Which one is real?: The problem of secondary mutation during mutagenesis in *Streptococcus pyogenes*, *Virulence*, 4(8), 1-2.

This Editorial is brought to you for free and open access by the Biochemistry and Molecular Medicine at Health Sciences Research Commons. It has been accepted for inclusion in Biochemistry and Molecular Medicine Faculty Publications by an authorized administrator of Health Sciences Research Commons. For more information, please contact hsrc@gwu.edu.

The prince and the pauper: Which one is real?

The problem of secondary mutation during mutagenesis in *Streptococcus pyogenes*

Kyu Hong Cho^{1,*}, Jordan Wright¹, Juan Svencionis¹, and Jeong-Ho Kim²

¹Department of Microbiology; Southern Illinois University; Carbondale, IL USA; ²Department of Biochemistry and Molecular Medicine; The George Washington University Medical Center; Washington, DC USA

Keywords: *Streptococcus pyogenes*, in vitro passages, secondary mutation, M protein, the *sagA/pel* locus

It has been known in the community studying the pathogenesis of *Streptococcus pyogenes* that spontaneous mutation occurs during passages in laboratory media or in animals, so some of the progeny's virulence can be changed, mainly due to the change of the expression of virulence factors.¹⁻³ This phenomenon is not only limited to *S. pyogenes*, but also happens to other pathogens.⁴⁻⁶ Like other pathogens, *S. pyogenes*, a strict human pathogen, produces virulence factors to attach to and invade host tissues, avoid the immune system, or degrade or kill host tissues and cells. These virulence factors are necessary to colonize, invade, and spread in the host. However, some of these factors are not necessary to grow in laboratory media, so reduced expression of these virulence factors could increase the fitness for growth in laboratory media. A notorious *S. pyogenes* virulence factor whose expression changes during passages in the laboratory is M protein. M protein is a cell wall-anchored protein with a coiled-coil structure and its important role in *S. pyogenes* pathogenesis has long been recognized as an adhesin and antiphagocytic factor.⁷ It is also the causative molecule of the streptococcal rheumatic heart disease. The antibodies generated against M protein cross react with tissues in the heart valves.⁸ Thus, when *S. pyogenes* loses M protein, its virulence is dramatically attenuated in animal infection models.

It has been a puzzle in the study of *S. pyogenes* whether or not the *sagA/pel* locus influences the expression of the M protein gene (*emm*). Several studies on the *sagA/pel* locus produced conflicting results on the regulation of M protein gene expression. However, the recent study by Zhou et al. in this issue of *Virulence* shows that the *sagA/pel* locus does not regulate the expression of M protein at least in the M1 strains that most studies for the *sagA/pel* locus used, and the confusion has been caused by secondary spurious mutations during mutagenesis.⁹ This study by Zhou et al. reminds us of the importance of awareness of spontaneous mutation or phase variation possibilities during mutagenesis of *S. pyogenes*. During laboratory passages, the expression of other virulence

factors such as capsule and the protease SpeB changes,³ so this phenomenon seems not to be limited to M protein.

Recent genomic studies show that genetic changes in *S. pyogenes* occur not only during passages in the laboratory (in vitro passage) but also during infection (in vivo passage).¹⁰⁻¹³ Interestingly, these genetic changes are selected mostly in the virulence factor-regulator genes of *mga*, *covRS*, and *ropB*. In the study by Zhou et al., the expression of both genes of M protein and its regulator Mga decreased in strains producing low levels of M protein after lab passages, and there was no mutation in the *emm* and *mga* promoter region. However, since the expression of *mga* is autoregulated,¹⁴ it cannot be ruled out that Mga was mutated in the study. The major virulence factors regulated by Mga, CovRS, and RopB are M protein, capsule, and SpeB, which are the same virulence factors whose expression was changed after laboratory passages. Therefore, the selected mutations during in vitro passages might also be concentrated in these regulators.

To prevent improper interpretation of a function attributed to a gene due to spurious secondary mutations, mutagenesis should be carefully performed and the phenotype of mutants should be carefully examined. *S. pyogenes* starts to die quickly after reaching the stationary phase in THY medium (most popular lab medium for *S. pyogenes*) probably due to the lack of nutrients, low pH (~pH 5.5) etc. It has been known that cells go through more mutation during conditions not favorable for growth such as in the stationary phase.^{15,16} Thus, during mutagenesis, cells should not stay in the stationary phase too long. After mutagenesis, the phenotype of several mutant colonies, not just one or two, should be examined. The expression of the virulence factors M protein, capsule, and SpeB can be easily examined phenotypically. The production level of capsule is easily determined by the appearance and size of colonies on agar plates. The expression level of the major secreted protease SpeB can be examined by plating cells on skim milk agar plates.¹⁷ Most, if not all, *S. pyogenes* strains form aggregates during stationary phase in liquid media, and M protein is involved in the aggregate formation through

*Correspondence to: Kyu Hong Cho; Email: kyuhongcho@gmail.com

Submitted: 10/08/2013; Accepted: 10/09/2013

<http://dx.doi.org/10.4161/viru.26767>

Comment on: Zhou Y, Hanks TS, Feng W, Li J, Liu G, Liu M, Lei B. The *sagA/pel* locus does not regulate the expression of the M protein of the M1T1 lineage of group A Streptococcus. *Virulence* 2013; 4:698–706; PMID: 24121654; <http://dx.doi.org/10.4161/viru.26413>

homophilic protein–protein interactions.¹⁸ However, other cell surface proteins such as M-like proteins, Protein H, and Protein F are also responsible for the aggregation. Thus, it is wise not to pick non-aggregated colonies if other colonies are aggregated in liquid media. However, when the expression of M protein needs to be tested, a direct method such as western blotting should be performed.

Another way to confirm the function of a gene attributed after mutagenesis is to perform the complementation test. Most commonly, multicopy plasmids are used to complement a gene or genes in *S. pyogenes*. Complementation with a multicopy plasmid is easy to perform, and if it works, the gene can be manipulated easily by deleting a motif or changing key amino acids through inverse PCR. However, complementation with a multicopy plasmid does not work in some cases, perhaps due to the issues

of the level or the location of expression. Another way to perform complementation is to insert the same gene using a suicide vector into the mutagenized site. If an intact gene is introduced into the original site in the chromosome, then the problems of copy number and expression location can be solved. Using this method, we succeeded in complementing genes whose complementation with a multicopy plasmid did not work.^{17,19} Previously, McShan et al. developed a streptococcal suicide vector system that is capable of site-specific integration.²⁰ This vector system contains *S. pyogenes* phage genes to facilitate the vector's integration into a serine tRNA gene, so this system can also be a useful system for complementation with a single copy gene.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

- Engleberg NC, Heath A, Miller A, Rivera C, DiRita VJ. Spontaneous mutations in the CsrRS two-component regulatory system of *Streptococcus pyogenes* result in enhanced virulence in a murine model of skin and soft tissue infection. *J Infect Dis* 2001; 183:1043-54; PMID:11237829; <http://dx.doi.org/10.1086/319291>
- Simpson WJ, Cleary PP. Expression of M type 12 protein by a group A streptococcus exhibits phaselike variation: evidence for coregulation of colony opacity determinants and M protein. *Infect Immun* 1987; 55:2448-55; PMID:2443452
- Leonard BA, Woischnik M, Podbielski A. Production of stabilized virulence factor-negative variants by group A streptococci during stationary phase. *Infect Immun* 1998; 66:3841-7; PMID:9673270
- Brummer E, Restrepo A, Hanson LH, Stevens DA. Virulence of *Paracoccidioides brasiliensis*: the influence of in vitro passage and storage. *Mycopathologia* 1990; 109:13-7; PMID:2139178; <http://dx.doi.org/10.1007/BF00437001>
- Nischik N, Schade B, Dytnerka K, Długońska H, Reichmann G, Fischer HG. Attenuation of mouse-virulent *Toxoplasma gondii* parasites is associated with a decrease in interleukin-12-inducing tachyzoite activity and reduced expression of actin, catalase and excretory proteins. *Microbes Infect* 2001; 3:689-99; PMID:11489417; [http://dx.doi.org/10.1016/S1286-4579\(01\)01425-3](http://dx.doi.org/10.1016/S1286-4579(01)01425-3)
- Werner JA, Kasten RW, Feng S, Sykes JE, Hodzic E, Salemi MR, Barthold SW, Chomel BB. Experimental infection of domestic cats with passaged genotype I *Bartonella henselae*. *Vet Microbiol* 2007; 122:290-7; PMID:17321078; <http://dx.doi.org/10.1016/j.vetmic.2007.01.018>
- Metzgar D, Zampolli A. The M protein of group A Streptococcus is a key virulence factor and a clinically relevant strain identification marker. *Virulence* 2011; 2:402-12; PMID:21852752; <http://dx.doi.org/10.4161/viru.2.5.16342>
- Nitsche-Schmitz DP, Chhatwal GS. Host-pathogen interactions in streptococcal immune sequelae. *Curr Top Microbiol Immunol* 2013; 368:155-71; PMID:23212184; http://dx.doi.org/10.1007/82_2012_296
- Zhou Y, Hanks TS, Feng W, Li J, Liu G, Liu M, Lei B. The *sagA/pel* locus does not regulate the expression of the M protein of the MIT1 lineage of Group A Streptococcus. *Virulence* 2013; 4:698-706; PMID:24121654 <http://dx.doi.org/10.4161/viru.26413>
- Carroll RK, Shelburne SA 3rd, Olsen RJ, Suber B, Sahasrabhojane P, Kumaraswami M, Beres SB, Shea PR, Flores AR, Musser JM. Naturally occurring single amino acid replacements in a regulatory protein alter streptococcal gene expression and virulence in mice. *J Clin Invest* 2011; 121:1956-68; PMID:21490401; <http://dx.doi.org/10.1172/JCI45169>
- Horstmann N, Sahasrabhojane P, Suber B, Kumaraswami M, Olsen RJ, Flores A, Musser JM, Brennan RG, Shelburne SA 3rd. Distinct single amino acid replacements in the control of virulence regulator protein differentially impact streptococcal pathogenesis. *PLoS Pathog* 2011; 7:e1002311; PMID:22028655; <http://dx.doi.org/10.1371/journal.ppat.1002311>
- Ikebe T, Ato M, Matsumura T, Hasegawa H, Sata T, Kobayashi K, Watanabe H. Highly frequent mutations in negative regulators of multiple virulence genes in group A streptococcal toxic shock syndrome isolates. *PLoS Pathog* 2010; 6:e1000832; PMID:20368967; <http://dx.doi.org/10.1371/journal.ppat.1000832>
- Vahling CM, McIver KS. Identification of residues responsible for the defective virulence gene regulator Mga produced by a natural mutant of *Streptococcus pyogenes*. *J Bacteriol* 2005; 187:5955-66; PMID:16109937; <http://dx.doi.org/10.1128/JB.187.17.5955-5966.2005>
- Okada N, Geist RT, Caparon MG. Positive transcriptional control of *mry* regulates virulence in the group A streptococcus. *Mol Microbiol* 1993; 7:893-903; PMID:8483419; <http://dx.doi.org/10.1111/j.1365-2958.1993.tb01180.x>
- Kivisaar M. Stationary phase mutagenesis: mechanisms that accelerate adaptation of microbial populations under environmental stress. *Environ Microbiol* 2003; 5:814-27; PMID:14510835; <http://dx.doi.org/10.1046/j.1462-2920.2003.00488.x>
- Kivisaar M. Mechanisms of stationary-phase mutagenesis in bacteria: mutational processes in pseudomonads. *FEMS Microbiol Lett* 2010; 312:1-14; PMID:20572869; <http://dx.doi.org/10.1111/j.1574-6968.2010.02027.x>
- Cho KH, Caparon MG. tRNA modification by GidA/MnmE is necessary for *Streptococcus pyogenes* virulence: a new strategy to make live attenuated strains. *Infect Immun* 2008; 76:3176-86; PMID:18426891; <http://dx.doi.org/10.1128/IAI.01721-07>
- Frick IM, Mörgelin M, Björck L. Virulent aggregates of *Streptococcus pyogenes* are generated by homophilic protein-protein interactions. *Mol Microbiol* 2000; 37:1232-47; PMID:10972839; <http://dx.doi.org/10.1046/j.1365-2958.2000.02084.x>
- Kang SO, Caparon MG, Cho KH. Virulence gene regulation by CvfA, a putative RNase: the CvfA-enolase complex in *Streptococcus pyogenes* links nutritional stress, growth-phase control, and virulence gene expression. *Infect Immun* 2010; 78:2754-67; PMID:20385762; <http://dx.doi.org/10.1128/IAI.01370-09>
- McShan WM, Tang YF, Ferretti JJ. Bacteriophage T12 of *Streptococcus pyogenes* integrates into the gene encoding a serine tRNA. *Mol Microbiol* 1997; 23:719-28; PMID:9157243; <http://dx.doi.org/10.1046/j.1365-2958.1997.2591616.x>