

8-22-2017

# Anti-HERV-K (HML-2) Capsid Antibody Responses in HIV Elite Controllers.

Miguel de Mulder

*George Washington University*

Devi SenGupta


Steven G Deeks

Jeffrey N Martin

Christopher D Pilcher

*See next page for additional authors*

Follow this and additional works at: [https://hsrc.himmelfarb.gwu.edu/smhs\\_microbio\\_facpubs](https://hsrc.himmelfarb.gwu.edu/smhs_microbio_facpubs)

 Part of the [Medical Immunology Commons](#), [Medical Microbiology Commons](#), [Tropical Medicine Commons](#), [Virology Commons](#), and the [Virus Diseases Commons](#)

## APA Citation

de Mulder, M., SenGupta, D., Deeks, S., Martin, J., Pilcher, C., Hecht, F., Sacha, J., Nixon, D., & Michaud, H. (2017). Anti-HERV-K (HML-2) Capsid Antibody Responses in HIV Elite Controllers.. *Retrovirology* [electronic resource], 14 (1). <http://dx.doi.org/10.1186/s12977-017-0365-2>

This Journal Article is brought to you for free and open access by the Microbiology, Immunology, and Tropical Medicine at Health Sciences Research Commons. It has been accepted for inclusion in Microbiology, Immunology, and Tropical Medicine Faculty Publications by an authorized administrator of Health Sciences Research Commons. For more information, please contact [hsrc@gwu.edu](mailto:hsrc@gwu.edu).

---

**Authors**

Miguel de Mulder, Devi SenGupta, Steven G Deeks, Jeffrey N Martin, Christopher D Pilcher, Frederick M Hecht, Jonah B Sacha, Douglas F Nixon, and Henri-Alexandre Michaud

RESEARCH

Open Access



# Anti-HERV-K (HML-2) capsid antibody responses in HIV elite controllers

Miguel de Mulder<sup>1\*</sup> , Devi SenGupta<sup>2,3</sup>, Steven G. Deeks<sup>4</sup>, Jeffrey N. Martin<sup>5</sup>, Christopher D. Pilcher<sup>4</sup>, Frederick M. Hecht<sup>4</sup>, Jonah B. Sacha<sup>6</sup>, Douglas F. Nixon<sup>1,3</sup> and Henri-Alexandre Michaud<sup>3,7</sup>

## Abstract

**Background:** Human endogenous retroviruses (HERVs) comprise approximately 8% of the human genome and while the majority are transcriptionally silent, the most recently integrated HERV, HERV-K (HML-2), remains active. During HIV infection, HERV-K (HML-2) specific mRNA transcripts and viral proteins can be detected. In this study, we aimed to understand the antibody response against HERV-K (HML-2) Gag in the context of HIV-1 infection.

**Results:** We developed an ELISA assay using either recombinant protein or 164 redundant “15mer” HERV-K (HML-2) Gag peptides to test sera for antibody reactivity. We identified a total of eight potential HERV-K (HML-2) Gag immunogenic domains: two on the matrix (peptides 16 and 31), one on p15 (peptide 85), three on the capsid (peptides 81, 97 and 117), one on the nucleocapsid (peptide 137) and one on the QP1 protein (peptide 157). Four epitopes (peptides 16, 31, 85 and 137) were highly immunogenic. No significant differences in antibody responses were found between HIV infected participants ( $n = 40$ ) and uninfected donors ( $n = 40$ ) for 6 out of the 8 epitopes tested. The antibody response against nucleocapsid (peptide 137) was significantly lower ( $p < 0.001$ ), and the response to QP1 (peptide 157) significantly higher ( $p < 0.05$ ) in HIV-infected adults compared to uninfected individuals. Among those with HIV infection, the level of response against p15 protein (peptide 85) was significantly lower in untreated individuals controlling HIV (“elite” controllers) compared to untreated non-controllers ( $p < 0.05$ ) and uninfected donors ( $p < 0.05$ ). In contrast, the response against the capsid protein (epitopes 81 and 117) was significantly higher in controllers compared to uninfected donors ( $p < 0.001$  and  $< 0.05$  respectively) and non-controllers ( $p < 0.01$  and  $< 0.05$ ). Peripheral blood mononuclear cells (PBMCs) from study participants were tested for responses against HERV-K (HML-2) capsid recombinant peptide in gamma interferon (IFN- $\gamma$ ) enzyme immunospot (Elispot) assays. We found that the HERV-K (HML-2) Gag antibody and T cell response by Elispot were significantly correlated.

**Conclusions:** HIV elite controllers had a strong cellular and antibody response against HERV-K (HML-2) Gag directed mainly against the Capsid region. Collectively, these data suggest that anti-HERV-K (HML-2) antibodies targeting capsid could have an immunoprotective effect in HIV infection.

**Keywords:** HIV, HERV-K, Antibodies, Gag, Elite Controllers, Viremic non-controllers

## Background

Human endogenous retroviruses (HERVs) are fossil remnants of inherited retroviruses which were endogenized into the genome, and comprise about 5–8% of the human genome [1]. Their ability to replicate or produce

infectious particles is impaired by host restriction [2, 3] and they are now considered to be stably integrated, largely silent, and transmitted in a Mendelian fashion [4]. Three major HERV classes have been identified and classified according to their polymerase gene (*pol*) sequence homology with exogenous retroviruses. Class I, II and III HERVs have similarities with gammaretroviruses, betaretroviruses and spumaviruses, respectively [5]. To date, endogenous homologues to lentiviruses have not been described in the human genome.

\*Correspondence: mdemulder@gwu.edu

<sup>1</sup> Department of Microbiology, Immunology, and Tropical Medicine, The George Washington University, Ross Hall 604, 2300 Eye St. NW, Washington, DC 20037, USA

Full list of author information is available at the end of the article

HERV-K (HML-2), a class II HERV, with *gag*, *pro*, *pol* and *env* genes, flanked by two Long Terminal Repeats (LTR), is the most recently integrated into the genome and under certain circumstances can express proteins [6, 7]. HERV-K (HML-2) expression has been associated with some autoimmune diseases [8–13] and cancers [14–19], and mRNA transcripts and proteins can be found in tumor tissues. Translated HERV proteins can induce an immune response that correlates with disease progression or regression in some cancers [20–25].

We, and others, have previously shown that HERV-K (HML-2) can be reactivated in HIV infection [26–28]. The mechanisms leading to HERV-K (HML-2) expression are still being elucidated, but HIV Vif and Tat proteins have been implicated [27, 29]. However, it appears that the transactivation of HERV-K by exogenous HIV is more complex than initial studies suggested. In a previous study, we showed that HIV induced a skewed expression of HERV-K (HML-2) Env which favored the surface cell expression of the transmembrane envelope glycoprotein (TM) at the expense of the surface unit (SU). We showed that isolated HERV-K specific T-cell clones and HA137, a human anti-HERV-K (HML-2) TM antibody, eliminated HIV infected cells in vitro [26–28, 30, 31].


To further characterize the role of the anti-HERV-K (HML-2) immune response in HIV infection, we investigated the antibody response to HERV-K (HML-2) Gag in HIV infected participants. In this study, we showed that strong anti-HERV-K (HML-2) capsid response is more frequently found in elite controllers (ECs) compared to

viremic non-controllers (VNCs) and HIV-negative low risk donors (SNLR). This response correlated with the HERV-K (HML-2) capsid T cell response. We mapped the antibody response and characterized an antibody pattern signature in ECs that significantly differed from the ones found VNCs, suggesting that the anti-HERV-K (HML-2) antibody response could play a role in the control of infection.

## Results

### The anti-HERV-K (HML-2) Capsid response correlates with anti-HERV Gag T-cell response in elite controllers

We first evaluated the antibody response against HERV-K (HML-2) recombinant capsid protein in uninfected donors and in untreated HIV-infected participants who were categorized as ECs or VNCs (Fig. 1). Although no significant differences were found in the magnitude of the antibody response between HIV-infected adults and HIV-negative low risk donors (SNLR), when the HIV-infected cohort was classified according to clinical status, we found that ECs had significantly higher level of antibodies against HERV-K (HML-2) capsid compared to SNLR ( $p < 0.01$ , Kruskal–Wallis test) and VNC ( $p < 0.001$ , Kruskal–Wallis test) (Fig. 1a). Since 85% of ECs developed a moderate or a strong anti-HERV Gag B-cell response (compared to 15% for VNCs), we investigated whether ECs also had a T-cell response against HERV-K (HML-2) Gag. We found that the HERV-K (HML-2) Gag antibody and T-cell response by Elispot were significantly correlated ( $p = 0.0047$ , Spearman test) (Fig. 1b).



### Identification of linear antibody epitopes on HERV-K (HML-2) Gag proteins

To further characterize anti-HERV-K (HML-2) Gag responses during HIV infection we used a set of 164 redundant “15mer” peptides overlapping by 11 amino acids in a peptide-based ELISA assay to map immunogenic domains. We used sera from 8 SNLR and confirmed that HIV uninfected donors had a low basal level of antibodies against HERV-K (HML-2) Gag, as previously published [32]. Among all tested peptides, 8 sequences had significant differences with the basal level. We identified 4 sequences with a higher reactivity with human sera and 4 sequences associated with a lower reactivity (Table 1; Fig. 2; Additional file 1: Fig. S1). The epitopes are distributed on HERV-K (HML-2) Gag as follows: two epitopes on matrix (MA, peptides 16 and 31), one epitope on p15 (peptide 58), three epitopes on capsid (CA, peptides 81, 97 and 117), one epitope on nucleocapsid (NC, peptide 137) and one epitope on the QP1 protein (peptide 157). These results suggest that each protein domain has different antibody immunogenicity to HERV-K (HML-2) proteins (Fig. 2). Indeed, sera from SNLR participants strongly reacted to the nucleocapsid epitope but not to the capsid (Fig. 2). Thus, domains can be classified as poorly immunogenic, such as the capsid or the QP1 and QP2 proteins, or immunogenic such as matrix, p15 and nucleocapsid.


### Elite controllers and viremic non-controllers have distinct antibody patterns

We then used the eight peptide epitopes identified above to perform a serological screen on 40 HIV-infected and 40 SNLR sera samples (Fig. 3). The differences between these two groups were not significant for 6 out of 8 epitopes (Fig. 3). The responses against MA (peptides 16 and 31) and p15 (peptide 58) were either slightly decreased or not changed upon HIV infection (Fig. 3a–c). However, the response against the most immunogenic domain (peptide 137) was significantly lower ( $p < 0.001$ ,  $q = 0.0004$ , Kruskal–Wallis test) in HIV infected subjects


**Table 1** Sequence identification of HERV-K (HML-2) Gag epitopes

Epitope	Sequence	Protein	Immunogenicity
16	KRIGKELKQAGRKGN	MA	Medium
31	KKSQKETESLHCEYV	MA	Medium
58	GYPGMPPAPQGRAPY	P15	Medium
81	GVKQYGPNSPYMRTL	CA	Low
97	NPPVNIDADQLLGIG	CA	Low
117	SIADKARKVIVELM	CA	Low
137	KCYNCGQIGHLKKNC	NC	High
157	PIQPFVPPQGFQGGQP	QP1	Low

MA matrix, CA capsid, NC nucleocapsid



**Fig. 2** Antibody mapping of anti-HERV-K (HML-2) responses in healthy donors. Sera from 8 seronegative low risk healthy donors (SNLR) were used for antibody epitope identification by ELISA. The 7 sub-units: matrix (MA), SP1, p15, capsid (CA), nucleocapsid (NC) and QP 1 and 2 are represented by 164 redundant 15mers named by their number corresponding to their rank in the list. The lines represent the average of the 8 individuals and duplicate signals (OD). Background was determined by the average of OD from each peptide. The statistical significance between the different groups was established using a Kruskal–Wallis and Dunn’s Multiple Comparison test and a  $p$  value  $< 0.05$  was considered as significant. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ ; \*\*\*\* $p < 0.0001$





(Fig. 3g). Although the response against peptide 97 did not differ based on HIV status, the response against the two other epitopes present on the capsid, peptide 81 and 117, trended towards being higher in those infected with HIV ( $p = 0.05$ ,  $q = 0.07$  and  $p = 0.08$ ,  $q = 0.1$  respectively, Kruskal–Wallis test) (Fig. 3d–f). The response against peptide 157 from the QP1 protein was the only that was significantly higher among those with HIV infection ( $p < 0.05$ ,  $q = 0.01$  Kruskal–Wallis test) (Fig. 3h).

To better understand the potential role of these antibody responses in HIV infection, we categorized the cohort based on clinical status. We used sera samples from 20 elite controllers (ECs) and 20 viremic non-controllers (VNCs). No differences were observed for the responses against peptides 16, 31, 97, 137 and 157 (Fig. 4a, b, e, g, h). However, significant differences were observed for the response against p15 (peptide 58). The response was significantly lower in elite controllers compared to SNLR ( $p < 0.05$ , Kruskal–Wallis test) and VNCs ( $p < 0.05$ , Kruskal–Wallis test), while there was no difference between VNCs and SNLR (Fig. 4c). In

contrast, antibody responses against capsid epitopes 81 and 117 were significantly higher in ECs compared to SNLR ( $p < 0.001$  and  $< 0.05$ , respectively, Kruskal–Wallis test) and VNCs ( $p < 0.01$  and  $< 0.05$ , Kruskal–Wallis test), but no differences were detected between VNCs and SNLR (Fig. 4f). We found that 90% of ECs had a moderate or a strong response against peptide 81 (25% for VNCs). Furthermore, we found a positive correlation for responses against capsid epitopes 81 and 117 ( $p = 0.0122$ ,  $r = 0.5768$ , Spearman test) in ECs that was not found in VNCs (Fig. 5a). However, compared to the response against peptide 58, we found a trend towards an inverse relationship between the antibody responses against HERV-K (HML-2) capsid and p15 developed in VNCs. Collectively, the data show that VNCs and ECs developed different anti-HERV-K (HML-2) gag antibody responses (Fig. 5b, c).


When analyzing the different groups according to their clinical status we found that VNC had low levels of anti-capsid antibodies, and there was a significant inverse correlation between the anti peptide 81 or anti-HERV-K



(HML-2) recombinant capsid responses and HIV viremia ( $p = -0.4879$ ,  $r = 0.0291$  and  $p = 0.0056$ ,  $r = -0.5955$  respectively, Spearman test) (Fig. 6). However, no correlation between CD4+T cell counts and anti-HERV-K (HML-2) Gag responses were detected, either in ECs or in VNCs (data not shown).

### Discussion

We, and others, have previously shown that HIV infection reactivates HERV-K (HML-2), leading to HERV-K (HML-2) Gag and Env protein production [27, 33]. In this study, we hypothesized that HIV infection leads to a HERV-K (HML-2) capsid antibody response. We focused



**Fig. 6** Correlations between HIV viremia and anti-HERV-K (HML-2) capsid responses. The correlation between the anti-HERV-K (HML-2) capsid response and viral load was determined by comparing the responses against peptide 81 (a) and the recombinant HERV-K (HML-2) capsid protein (b) measured by peptide-based ELISA for 20 viremic non-controllers. The statistical significance between the different responses was established using the non-parametric Spearman test. Figure 6a shows a representative result of four independent experiments. Figure 6b shows the mean of four independent experiments. A  $p$  value  $<0.05$  was considered as significant.  $*p < 0.05$ ;  $**p < 0.01$

on responses to HERV-K (HML-2) capsid based on our previous findings that elite controllers (EC) developed an anti-HERV-K (HML-2) capsid cellular immune response [30]. We show here that a strong antibody response against HERV-K (HML-2) capsid can be detected in ECs. To better characterize the anti-HERV-K (HML-2) response, we mapped responses to linear epitopes on the full HERV-K (HML-2) Gag sequence. Responses against HERV-K (HML-2) Gag were not changed by HIV infection, except for responses to an epitope present on the QP1 protein (epitope 157), which were higher in HIV-1 infected participants compared to controls. The antibody response against nucleocapsid (epitope 137) was significantly lower in HIV infected participants compared to uninfected individuals.

When we compared the anti-HERV-K (HML-2) gag antibody response between ECs and VNCs, we saw a distinct antibody pattern, characterized by a significantly higher anti-capsid antibodies (specific to peptides 81 and 117) and lower anti-p15 antibodies (epitope 58) in controllers. The plasma HIV RNA levels were strongly inversely correlated in VNCs who had an anti-HERV-K (HML-2) capsid response.

It has been previously established that HERV-K (HML-2) expression is tightly associated with HIV viral transcription and activity *in vitro* and *in vivo* [31, 34–37]. However, ECs have limited HIV replication activity, suggesting that the induction of the anti HERV-K (HML-2) capsid response in ECs is not caused by HIV-induced HERV-K capsid expression, as has been previously described for the HERV-K (HML-2) envelope antibody response [31, 36, 37]. However, a longitudinal study would be more informative to determine potential causality.

In a previous report, we showed that expression of HERV-K Env proteins following HIV infection is skewed towards a predominant expression of the HERV-K transmembrane protein compared to the surface unit protein [31]. This suggested an HIV/HERV-K (HML-2) interaction far more complex than previous studies have proposed. In the case of HERV-K (HML-2) Gag expression, antibody profiles found in VNCs and ECs seem to reinforce this complexity. Anti-HERV-K (HML-2) capsid responses in VNCs are not significantly different than those found in controls, but they seem to be lower in early and late stages of HIV infection despite an increase of HERV-K (HML-2) expression. Patients who naturally control HIV infection are more likely to have a strong antibody response against HERV-K capsid, but the antibody response against p15 (peptide 58) was strongly decreased in ECs compared to VNCs and SNLR. Further longitudinal studies are needed to understand the chronology and the cause of this dichotomy.

A second objective of our study was to characterize the role of anti HERV-K (HML-2) capsid responses in ECs. Antibodies against viral gag proteins are not unexpected in HIV patients [38] and the anti-HIV p24 response correlates with control of disease progression [38]. Antibodies against capsid could bind cells resulting in their lysis and formation of immunocomplexes. Those immunocomplexes might interact with innate immune cells such as NK cells, macrophages or dendritic cells and promote their activation and induction of a cellular immune response [39, 40]. This may explain why both T and B cell responses directed against HIV p24 correlated with the status of controllers. A similar hypothesis could be applied for anti-HERV-K (HML-2) capsid antibodies,



with EC having both an anti-HIV p24 and an anti-HERV-K (HML-2) capsid response as a way to reinforce their antiviral response.

## Conclusion

In this study, we identified linear immunogenic antibody epitopes on HERV-K (HML-2) gag proteins. We found that elite controllers had a distinctive antibody pattern compared to viremic non-controllers and HIV seronegative participants. Although further studies are needed to elucidate how these responses could be involved in the control of viremia, it reinforces the importance of studying HERV-K (HML-2) capsid immune responses in HIV infection.

## Methods

### Study populations

Samples of peripheral blood mononuclear cells (PBMCs) were selected from participants in a San Francisco-based HIV-infected cohort: OPTIONS (n = 40). Samples from HIV-negative controls were obtained from individuals who donated blood to the Stanford blood bank. Studies were performed on cryopreserved PBMCs and sera.

PBMC and sera samples were obtained from the following categories of chronically HIV-infected individuals: 20 elite controllers (EC: naive for treatment, undetectable viral load for two years, CD4 > 350) and 20 untreated virologic non-controllers (VNC; naive for treatment, viral load >2000 copies/mL).

### ELISA

A set of 164 overlapping “15-mer” HERV-K (HML-2) Gag peptides (JPT Peptide Technologies, Berlin, Germany), based on HERV-K102 sequence (AF164610), were used to comprehensively map the HERV-K (HML-2) antibody response (Additional file 2: Fig. S2). Positive signals were confirmed by peptides produced by two other companies (New England Peptide and Gene Script). 96 microtiter wells plate (Nunc-Immuno Plate MaxiSorp Surface) were coated for 1 h at 37 °C with peptides at 10 µg/ml in PBS or over-night at 4 °C with recombinant protein (GeneArt) at 5 µg/ml in PBS. Plates were then washed 3 times with 200 µL of PBS/0.05%-Tween 20 and blocked with 100 µL of blocking buffer (PBS/2.5%-BSA) at room temperature (RT). The samples were diluted in blocking buffer and incubated 2 h at RT in duplicates. Plates were then washed 3 times with 200 µL of PBS/0.05%-Tween 20. An anti-human IgG or anti-human IgM HRP-conjugated secondary antibody was diluted at 1:1000 in blocking buffer and incubated at RT for 1 h. Plates were then washed 6 times with 200 µL of PBS/0.05%-Tween 20 and incubated for 10 min with 100 µL of TMB (Invitrogen). Addition of 50 µL H<sub>2</sub>SO<sub>4</sub> 2 M stopped the reaction. The

plates were read at 450 and 690 nm for the background on a plate reader. Background from 690 nm uncoated wells and twice the background from 450 nm PBS-BSA (negative control) were subtracted from the mean absorbance of the coated wells (corrected OD). For the detection of anti-Gag antibodies, sera were used at 1:400. ODs were normalized with serum from a high responder in a standard curve. The STDEV intra experiment was less than 4%.

For the response against the recombinant HERV-K (HML-2) capsid and the peptide 81, we defined the humoral responses for elite controllers as followed: the response was considered moderate if the corrected OD is greater than the mean of corrected OD for SNLR and strong if the corrected OD is greater than twice the mean of corrected OD for SNLR.

### ELISPOT assays

The ELISPOT assay has been described previously [30]. In brief, 96-well plates (Millipore, Billerica, MA) were coated with human monoclonal anti-interferon gamma (IFN-γ) immunoglobulin (Mabtech, Mariemont, OH). After plates were washed and blocked with 10% fetal calf serum, PBMCs were added at a concentration of 10<sup>5</sup> cells per well. Duplicate wells were prepared for each experimental condition. Spot totals for duplicate wells were averaged, and all spot numbers were normalized to numbers of (IFN-γ) spot forming units (SFU) per million PBMCs (SPM). The spot values from medium control wells were subtracted, after which a positive response to a peptide was defined as 50 SPM and 2 times the medium control value. The total magnitude of the HERV T cell response was calculated by adding up all of the individual peptide SPM values.

### Statistical analyses

To assess the distribution of the humoral responses obtained in this study, we used the D’Agostino and Pearson normality test. The results concluded that the populations were not normally distributed. According to this statement, we used non-parametric statistical tests to compare the humoral responses assayed by ELISA for each group. Multiple comparisons were performed in the 3 groups (SNLR, EC and VNC) with the Kruskal–Wallis and Dunn’s multiple comparison test for Fig. 4. Spearman correlation analyses were used to measure associations between different humoral responses and HIV viral load or CD4+T cells count for Figs. 1b, 5, and 6. The two-tailed Mann–Whitney *u* test was used to compare the humoral responses between HIV-1pos and HIV-1neg (SNLR) groups for Fig. 3. A *q* value was calculated using the original method of Benjamini and Hochberg and added on the figure when it was relevant. All tests were

conducted using GraphPad Prism, version 6.00 (GraphPad Software, San Diego, CA), with the statistical significance of the findings set at a *p* value of less than 0.05.

## Additional files

**Additional file 1: Fig. S1.** Antibody mapping of anti-HERV-K (HML-2) responses in study participants groups. Sera from Healthy donors, Elite Controllers and Viremic non-controllers were used for antibody epitope identification by ELISA. The 7 sub-units: matrix (MA), SP1, p15, capsid (CA), nucleocapsid (NC) and QP 1 and 2 are represented by 164 redundant 15mers named by their number corresponding to their rank in the list. The lines represent the average of the 8 individuals and duplicate signals (OD). Background was determined by the average of OD from each peptide. Peptides giving a signal significantly decreased are symbolized in grey and peptides giving a signal significantly increased are symbolized in black.

**Additional file 2: Fig. S2.** Strategy for designing overlapping peptides.

## Authors' contributions

MMR designed and performed the experiments and wrote the paper, DS helped design and perform the experiments, SGD helped design the experiments and provided clinical samples, JNM helped design the experiments and provided clinical samples, CDP provided clinical samples, FMH provided clinical samples, RASR helped design the experiments and co-wrote the paper, JBS helped design the experiments and co-wrote the paper, DFN helped design the experiments and co-wrote the paper, H-AM designed and performed the experiments and wrote the paper. All authors helped edit the paper. All authors read and approved the final manuscript.

## Author details

<sup>1</sup> Department of Microbiology, Immunology, and Tropical Medicine, The George Washington University, Ross Hall 604, 2300 Eye St. NW, Washington, DC 20037, USA. <sup>2</sup> Gilead Sciences Inc., Foster City, CA, USA. <sup>3</sup> Division of Experimental Medicine, Department of Medicine, University of California, San Francisco, San Francisco, CA, USA. <sup>4</sup> HIV/AIDS Program, Department of Medicine, University of California, San Francisco, San Francisco, CA, USA. <sup>5</sup> Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, CA, USA. <sup>6</sup> Division of Pathobiology and Immunology, Oregon Health and Science University, Portland, CA, USA. <sup>7</sup> Equipe Immunité et Cancer, Institut de Recherche en Cancérologie de Montpellier, Montpellier, France.

## Acknowledgements

We thank Christopher Ormsby, Mario Ostrowski, Dominic Paquin-Proulx and Greta Beckerle for helpful discussions. We thank Rasha Boulos and Sabine Chabaliel from the bioinformatics department of the Institut de Recherche en Cancérologie de Montpellier. We gratefully acknowledge reagents supplied from the National Institutes of Health AIDS Research and Reference Reagent Program.

## Competing interests

The authors declare that they have no competing interests.

## Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

The study was approved by the local Institutional Review Board (University of California San Francisco Committee on Human Research), research conducted according to the Declaration of Helsinki, and individuals gave written informed consent.

## Funding

We gratefully acknowledge reagents supplied from the National Institutes of Health (NIH) AIDS Research and Reference Reagent Program. This work was supported in part by grant number 108248-51-RGRL from amfAR, The Foundation for AIDS Research, The Bill and Melinda Gates Foundation (BMGF:01434000363), the Peter and Shelagh Godsoe Family Foundation through the AIDS Research Institute at UCSF, Pfizer Inc. through a sponsored research agreement, and funds from the NIH (R01 AI076059, R56 AI084113, and P51 OD011092). This research was supported in part by a grant from the NIH, University of California, San Francisco-Gladstone Institute of Virology and Immunology Center for AIDS Research (P30 AI027763) and the District of Columbia Developmental Center for AIDS Research (P30 AI087714). The Options study is funded in part by National Institute of Allergy and Infectious Diseases (P01 AI071713). The SCOPE cohort is supported by the NIH (R01 AI087145, K24 AI069994, U19 AI0961090, UL1 RR024131, and R24 AI067039).

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 6 April 2017 Accepted: 13 August 2017

Published online: 22 August 2017

## References

- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing and analysis of the human genome. *Nature*. 2001;409(6822):860–921.
- Goff SP. Retrovirus restriction factors. *Mol Cell*. 2004;16(6):849–59.
- Lee YN, Bieniasz PD. Reconstitution of an infectious human endogenous retrovirus. *PLoS Pathog*. 2007;3(1):e10.
- Subramanian RP, Wildschutte JH, Russo C, Coffin JM. Identification, characterization, and comparative genomic distribution of the HERV-K (HML-2) group of human endogenous retroviruses. *Retrovirology*. 2011;8:90.
- Boeke JD, Stoye JP. Retrotransposons, endogenous retroviruses, and the evolution of retroelements. In: Coffin JM, Hughes SH, Varmus HE, editors. *Retroviruses*. New York: Cold Spring Harbor; 1997.
- Kurth R, Bannert N. Beneficial and detrimental effects of human endogenous retroviruses. *Int J Cancer*. 2010;126(2):306–14.
- Jha AR, Nixon DF, Rosenberg MG, Martin JN, Deeks SG, Hudson RR, et al. Human endogenous retrovirus K106 (HERV-K106) was infectious after the emergence of anatomically modern humans. *PLoS ONE*. 2011;6(5):e20234.
- Armitage AE, Katzourakis A, de Oliveira T, Welch JJ, Belshaw R, Bishop KN, et al. Conserved footprints of APOBEC3G on Hypermutated human immunodeficiency virus type 1 and human endogenous retrovirus HERV-K(HML2) sequences. *J Virol*. 2008;82(17):8743–61.
- Ehrlhardt S, Seifert M, Schneider J, Ojak A, Zang KD, Mehraein Y. Human endogenous retrovirus HERV-K(HML-2) Rec expression and transcriptional activities in normal and rheumatoid arthritis synovia. *J Rheumatol*. 2006;33(1):16–23.
- Herve CA, Lugli EB, Brand A, Griffiths DJ, Venables PJ. Autoantibodies to human endogenous retrovirus-K are frequently detected in health and disease and react with multiple epitopes. *Clin Exp Immunol*. 2002;128(1):75–82.
- International HIVCS, Pereyra F, Jia X, McLaren PJ, Telenti A, de Bakker PI, et al. The major genetic determinants of HIV-1 control affect HLA class I peptide presentation. *Science*. 2010;330(6010):1551–7.
- Lai OY, Chen H, Michaud HA, Hayashi G, Kuebler PJ, Hultman GK, et al. Protective effect of human endogenous retrovirus K dUTPase variants on psoriasis susceptibility. *J Invest Dermatol*. 2012;132(7):1833–40.
- Sicat J, Sutkowski N, Huber BT. Expression of human endogenous retrovirus HERV-K18 superantigen is elevated in juvenile rheumatoid arthritis. *J Rheumatol*. 2005;32(9):1821–31.
- Buscher K, Trefzer U, Hofmann M, Sterry W, Kurth R, Denner J. Expression of human endogenous retrovirus K in melanomas and melanoma cell lines. *Cancer Res*. 2005;65(10):4172–80.

15. Goering W, Ribarska T, Schulz WA. Selective changes of retroelement expression in human prostate cancer. *Carcinogenesis*. 2011;32(10):1484–92.
16. Ishida T, Obata Y, Ohara N, Matsushita H, Sato S, Uenaka A, et al. Identification of the HERV-K gag antigen in prostate cancer by SEREX using autologous patient serum and its immunogenicity. *Cancer Immun*. 2008;8:15.
17. Katoh I, Mirova A, Kurata S, Murakami Y, Horikawa K, Nakakuki N, et al. Activation of the long terminal repeat of human endogenous retrovirus K by melanoma-specific transcription factor MITF-M. *Neoplasia*. 2011;13(11):1081–92.
18. Schanab O, Humer J, Gleiss A, Mikula M, Sturlan S, Grunt S, et al. Expression of human endogenous retrovirus K is stimulated by ultraviolet radiation in melanoma. *Pigment Cell Melanoma Res*. 2011;24(4):656–65.
19. Wang-Johanning F, Frost AR, Jian B, Azerou R, Lu DW, Chen DT, et al. Detecting the expression of human endogenous retrovirus E envelope transcripts in human prostate adenocarcinoma. *Cancer*. 2003;98(1):187–97.
20. Boller K, Janssen O, Schuldes H, Tonjes RR, Kurth R. Characterization of the antibody response specific for the human endogenous retrovirus HTDV/HERV-K. *J Virol*. 1997;71(6):4581–8.
21. Hahn S, Ugurel S, Hanschmann KM, Strobel H, Tondera C, Schadendorf D, et al. Serological response to human endogenous retrovirus K in melanoma patients correlates with survival probability. *AIDS Res Hum Retroviruses*. 2008;24(5):717–23.
22. Rakoff-Nahoum S, Kuebler PJ, Heymann JJ, Sheehy ME, Ortiz GM, Ogg GS, et al. Detection of T lymphocytes specific for human endogenous retrovirus K (HERV-K) in patients with seminoma. *AIDS Res Hum Retroviruses*. 2006;22(1):52–6.
23. Schiavetti F, Thonnard J, Colau D, Boon T, Coulie PG. A human endogenous retroviral sequence encoding an antigen recognized on melanoma by cytolytic T lymphocytes. *Cancer Res*. 2002;62(19):5510–6.
24. Zhao J, Rycaj K, Geng S, Li M, Plummer JB, Yin B, et al. Expression of human endogenous retrovirus type K envelope protein is a novel candidate prognostic marker for human breast cancer. *Genes Cancer*. 2011;2(9):914–22.
25. Wang-Johanning F, Radvanyi L, Rycaj K, Plummer JB, Yan P, Sastry KJ, et al. Human endogenous retrovirus K triggers an antigen-specific immune response in breast cancer patients. *Cancer Res*. 2008;68(14):5869–77.
26. Garrison KE, Jones RB, Meiklejohn DA, Anwar N, Ndhlovu LC, Chapman JM, et al. T cell responses to human endogenous retroviruses in HIV-1 infection. *PLoS Pathog*. 2007;3(11):e165.
27. Jones RB, Garrison KE, Mujib S, Mihajlovic V, Aidarus N, Hunter DV, et al. HERV-K-specific T cells eliminate diverse HIV-1/2 and SIV primary isolates. *J Clin Invest*. 2012;122(12):4473–89.
28. Michaud HA, SenGupta D, de Mulder M, Deeks SG, Martin JN, Kobie JJ, et al. Cutting edge: an antibody recognizing ancestral endogenous virus glycoproteins mediates antibody-dependent cellular cytotoxicity on HIV-1-infected cells. *J Immunol*. 2014;193(4):1544–8.
29. Contreras-Galindo R, Kaplan MH, Contreras-Galindo AC, Gonzalez-Hernandez MJ, Ferlenghi I, Giusti F, et al. Characterization of human endogenous retroviral elements in the blood of HIV-1-infected individuals. *J Virol*. 2012;86(1):262–76.
30. SenGupta D, Tandon R, Vieira RG, Ndhlovu LC, Lown-Hecht R, Ormsby CE, et al. Strong human endogenous retrovirus-specific T cell responses are associated with control of HIV-1 in chronic infection. *J Virol*. 2011;85(14):6977–85.
31. Michaud HA, de Mulder M, SenGupta D, Deeks SG, Martin JN, Pilcher CD, et al. Trans-activation, post-transcriptional maturation, and induction of antibodies to HERV-K (HML-2) envelope transmembrane protein in HIV-1 infection. *Retrovirology*. 2014;11:10.
32. Reis BS, Jungbluth AA, Frosina D, Holz M, Ritter E, Nakayama E, et al. Prostate cancer progression correlates with increased humoral immune response to a human endogenous retrovirus GAG protein. *Clin Cancer Res*. 2013;19(22):6112–25.
33. Gonzalez-Hernandez MJ, Swanson MD, Contreras-Galindo R, Cookinham S, King SR, Noel RJ Jr, et al. Expression of human endogenous retrovirus type K (HML-2) is activated by the Tat protein of HIV-1. *J Virol*. 2012;86(15):7790–805.
34. Bhardwaj N, Maldarelli F, Mellors J, Coffin JM. HIV-1 infection leads to increased transcription of human endogenous retrovirus HERV-K (HML-2) proviruses in vivo but not to increased virion production. *J Virol*. 2014;88(19):11108–20.
35. Contreras-Galindo R, Gonzalez M, Almodovar-Camacho S, Gonzalez-Ramirez S, Lorenzo E, Yamamura Y. A new real-time-RT-PCR for quantitation of human endogenous retroviruses type K (HERV-K) RNA load in plasma samples: increased HERV-K RNA titers in HIV-1 patients with HAART non-suppressive regimens. *J Virol Methods*. 2006;136(1–2):51–7.
36. Contreras-Galindo R, Kaplan MH, Markovitz DM, Lorenzo E, Yamamura Y. Detection of HERV-K(HML-2) viral RNA in plasma of HIV type 1-infected individuals. *AIDS Res Hum Retroviruses*. 2006;22(10):979–84.
37. Contreras-Galindo R, Lopez P, Velez R, Yamamura Y. HIV-1 infection increases the expression of human endogenous retroviruses type K (HERV-K) in vitro. *AIDS Res Hum Retrovir*. 2007;23(1):116–22.
38. French MA, Tanaskovic S, Law MG, Lim A, Fernandez S, Ward LD, et al. Vaccine-induced IgG2 anti-HIV p24 is associated with control of HIV in patients with a 'high-affinity' FcγRIIIa genotype. *Aids*. 2010;24(13):1983–90.
39. French MA, Center RJ, Wilson KM, Fleyfel I, Fernandez S, Schorcht A, et al. Isotype-switched immunoglobulin G antibodies to HIV Gag proteins may provide alternative or additional immune responses to 'protective' human leukocyte antigen-B alleles in HIV controllers. *Aids*. 2013;27(4):519–28.
40. Michaud HA, Gomard T, Gros L, Thiolon K, Nasser R, Jacquet C, et al. A crucial role for infected-cell/antibody immune complexes in the enhancement of endogenous antiviral immunity by short passive immunotherapy. *PLoS Pathog*. 2010;6(6):e1000948.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

