

2014

Anesthesia for ambulatory surgery in a child with hyposensitivity to pain

Claude Abdallah

George Washington University

Follow this and additional works at: http://hsrc.himmelfarb.gwu.edu/smhs_anesth_facpubs



Part of the [Anesthesia and Analgesia Commons](#)

Recommended Citation

Abdallah, C. (2014). Anesthesia for ambulatory surgery in a child with hyposensitivity to pain. *World Journal of Anesthesiology*, 3(2), 189-190.

This Journal Article is brought to you for free and open access by the Anesthesiology and Critical Care Medicine at Health Sciences Research Commons. It has been accepted for inclusion in Anesthesiology and Critical Care Medicine Faculty Publications by an authorized administrator of Health Sciences Research Commons. For more information, please contact hsrc@gwu.edu.

EAnesthesia for ambulatory surgery in a child with hyposensitivity to pain

Claude Abdallah

CITATION	Abdallah C. Anesthesia for ambulatory surgery in a child with hyposensitivity to pain. <i>World J Anesthesiol</i> 2014; 3(2): 189-190
URL	http://www.wjgnet.com/2218-6182/full/v3/i2/189.htm
DOI	http://dx.doi.org/10.5313/wja.v3.i2.189
OPEN ACCESS	Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.
CORE TIP	Congenital hyposensitivity to pain is a condition with predisposition to injury. In patients with congenital hyposensitivity to pain/Hereditary sensory and autonomic neuropathy (HSAN), knowledge regarding anesthetic requirements and complications derives from individual case reports, or small case series. Different categories of HSAN have been described. Complications in the immediate perioperative period have been described such as mild hypothermia and cardiovascular events, mostly bradycardia and hypotension. The majority of patients with hyposensitivity to pain reported in the literature have received standard anesthesia for surgery. Immobilization, prevention of autonomic reflexes, anxiolysis, and sedation are equally important aspects of the anesthetic management in patients with hyposensitivity to pain.
KEY WORDS	General anesthesia; Child; Pain; Hyposensitivity; Surgery
COPYRIGHT	© 2014 Baishideng Publishing Group Co., Limited. All rights reserved.
COPYRIGHT LICENSE	Order reprints or request permissions: bpgoffice@wjgnet.com

NAME OF JOURNAL	<i>World Journal of Anesthesiology</i>
ISSN	2218-6182 (online)
PUBLISHER	Baishideng Publishing Group Co., Limited , Flat C, 23/F., Lucky Plaza, 315-321 Lockhart Road, Wan Chai, Hong Kong, China
WEBSITE	http://www.wjgnet.com

ESPS Manuscript NO: 9653

Columns: CASE REPORT

Anesthesia for ambulatory surgery in a child with hyposensitivity to pain

Claude Abdallah

Claude Abdallah, Division of Anesthesiology, Children's National Medical Center, The George Washington University Medical Center, Washington, DC 20010-2970, United States

Author contributions: Abdallah C cited above had complete contribution to conception, writing, and critically revising the intellectual content and approval of the published version of this report.

Correspondence to: Claude Abdallah, MD, MS, Division of Anesthesiology, Children's National Medical Center, The George Washington University Medical Center, 111 Michigan Avenue, N.W., Washington, DC 20010-2970, United States. cabdalla@cnmc.org

Telephone: +1-202-4762025 Fax: +1-202-4765999

Received: February 22, 2014 Revised: April 19, 2014 Accepted: May 16, 2014

Published online: July 27, 2014

Abstract

Congenital hyposensitivity to pain is a condition with predisposition to injury. In these patients, knowledge regarding anesthetic requirements and complications derives from individual case reports, or small case series. Different categories have been described. In patients with hyposensitivity to

pain, preventing and treating anxiety as well as insuring immobilization, avoidance of triggering of autonomic reflexes, and sedation are integral aspects for a safe and adequate anesthetic management.

© 2014 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: General anesthesia; Child; Pain; Hyposensitivity; Surgery

Core tip: Congenital hyposensitivity to pain is a condition with predisposition to injury. In patients with congenital hyposensitivity to pain/Hereditary sensory and autonomic neuropathy (HSAN), knowledge regarding anesthetic requirements and complications derives from individual case reports, or small case series. Different categories of HSAN have been described. Complications in the immediate perioperative period have been described such as mild hypothermia and cardiovascular events, mostly bradycardia and hypotension. The majority of patients with hyposensitivity to pain reported in the literature have received standard anesthesia for surgery. Immobilization, prevention of autonomic reflexes, anxiolysis, and sedation are equally important aspects of the anesthetic management in patients with hyposensitivity to pain.

Abdallah C. Anesthesia for ambulatory surgery in a child with hyposensitivity to pain. *World J Anesthesiol* 2014; 3(2): 189-190 Available from: URL: <http://www.wjgnet.com/2218-6182/full/v3/i2/189.htm> DOI: <http://dx.doi.org/10.5313/wja.v3.i2.189>

INTRODUCTION

Congenital hyposensitivity to pain is a condition with predisposition to injury, often associated with a delay and difficulty in diagnosis. Anesthesia care of these children may pose a challenge secondary to the rarity of the disease, the

presence of unclassified congenital variants of pain hyposensitivity, and the limited information regarding anesthetic management.

CASE REPORT

The following is a description of a case of a 2-year-old patient, female gender, with mild developmental delay, who presented as a same day case for ear surgery. The duration of the surgery was expected to be 90 min. During the process of the interview, the mother revealed that her daughter is insensitive to pain, no further details were available. Otherwise the review of systems was negative. Premedication with oral midazolam 0.5 mg/kg was administered. Patient received general anesthesia with endotracheal intubation [Sevoflurane mask induction in O₂/N₂O followed by propofol (3 mg/kg □)], and narcotics were titrated to maintain spontaneous ventilation for a total dose of Fentanyl of 1 mcg/kg. No fluctuations in temperature were noticed during the case. Vital signs were stable. There was no delay in emergence from general anesthesia. Recovery room stay was not prolonged and was uneventful. Patient was discharged the same day without further need for narcotics postoperatively.

DISCUSSION

In patients with congenital hyposensitivity to pain/hereditary sensory and autonomic neuropathy (HSAN), knowledge regarding anesthetic requirements and complications derives from individual case reports, or small case series. Different levels and modalities of autonomic dysfunction and sensory loss have been described^[1]. Some patients do have tactile hyperesthesia, or partially preserved nociception with sometimes preserved mechanoreceptor, cooling, and warming sensations. Five types of HSAN have been categorized. HSAN □ is inherited with autosomal dominance. The age of onset of HSAN □ is between the 2nd and 4th decade of life while the other types are autosomal recessive with an earlier age of onset, usually at birth in type □ (familial

dysautonomia) or in infancy. HSANs I, II and III usually present with a profound reduction of pain perception, while HSAN IV has as a milder manifestation. Patients with HSAN I's may have intact visceral and peritoneal pain sensation with profound dysautonomia. Thermal perception is severely impaired in all HSAN types. Mild hyperhidrosis is associated in HSAN type I, hypohidrosis in type II and III, while severe anhidrosis with recurrent episodes of severe hyperpyrexia is associated to HSAN type IV. Type IV may present with unaffected sensitivity to touch, pressure and vibration. Mutilations may be common in all types of HSAN. The requirements for volatile anesthetics have been described as being within the range of standard population. Intraoperative opioids dosage has been reported to be less than standard, if not negligible. Because some HSAN patients may have anhidrosis, intraoperative hyperthermia cases have been reported^[2]. There is no description of malignant hyperthermia in association with the different HSAN types. Patients usually do not require opioids postoperatively. Complications in the perioperative care have been described as cardiovascular, such as bradycardia and hypotension and mild hypothermia. A cardiac arrest following management of a patient with HSAN IV has been published^[3]. Although there is report of an adult patient diagnosed with profound congenital insensitivity to pain who has undergone a major orthopedic surgery without receiving general anesthesia and narcotics; in the literature, patients with hyposensitivity to pain, have been documented to receive standard anesthesia for surgery. In patients with hyposensitivity to pain, preventing and treating anxiety as well as insuring immobilization, avoidance of triggering of autonomic reflexes, and sedation are integral aspects for a safe and adequate anesthetic management.

COMMENTS

Case characteristics

Pediatric patient with hyposensitivity to pain.

Clinical diagnosis

Anesthesia management of a pediatric patient with hyposensitivity to pain.

Differential diagnosis

Different categories of hereditary sensory and autonomic neuropathy are described.

Experiences and lessons

Immobilization, prevention of autonomic reflexes, anxiolysis, and sedation are equally important aspects of the anesthetic management in patients with hyposensitivity to pain.

Peer review

The author described anesthesia management in a child with congenital hyposensitivity to pain. The diagnosis is based on the history and clinical findings. The paper is good for publication.

REFERENCES

- 1 **Weingarten TN**, Sprung J, Ackerman JD, Bojanic K, Watson JC, Dyck PJ. Anesthesia and patients with congenital hyposensitivity to pain. *Anesthesiology* 2006; **105**: 338-345 [PMID: 16871068 DOI: 10.1097/00000542-200608000-00017]
- 2 **Nolano M**, Crisci C, Santoro L, Barbieri F, Casale R, Kennedy WR, Wendelschafer-Crabb G, Provitera V, Di Lorenzo N, Caruso G. Absent innervation of skin and sweat glands in congenital insensitivity to pain with anhidrosis. *Clin Neurophysiol* 2000; **111**: 1596-1601 [PMID: 10964070 DOI: 10.1016/S1388-2457(00)00351-5]
- 3 **Ergül Y**, Ekici B, and Keskin S. Cardiac arrest after anesthetic management in a patient with hereditary sensory autonomic neuropathy type IV. *Saudi J Anaesth* 2011; **5**: 93–95 [DOI: 10.4103/1658-354X.76486]

P- Reviewer: Yang ZJ **S- Editor:** Song XX **L- Editor:** A **E- Editor:**
Wu HL

